

BMJ Open Seeking culturally safe care: a qualitative systematic review of the healthcare experiences of women and girls who have undergone female genital mutilation/cutting

Catrin Evans,¹ Ritah Tweheyo,¹ Julie McGarry,¹ Jeanette Eldridge,² Juliet Albert,³ Valentine Nkoyo,⁴ Gina Marie Awoko Higginbottom¹

To cite: Evans C, Tweheyo R, McGarry J, *et al.* Seeking culturally safe care: a qualitative systematic review of the healthcare experiences of women and girls who have undergone female genital mutilation/cutting. *BMJ Open* 2019;**9**:e027452. doi:10.1136/bmjopen-2018-027452

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-027452>).

Received 23 October 2018
Revised 15 March 2019
Accepted 15 April 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY. Published by BMJ.

¹School of Health Sciences, University of Nottingham, Nottingham, UK

²Research and Learning Services, School of Health Sciences, University of Nottingham, Nottingham, UK

³Department of Midwifery, Imperial College Healthcare NHS Trust, London, UK

⁴Mojatu Foundation, Nottingham, UK

Correspondence to

Dr Catrin Evans;
catrin.evans@nottingham.ac.uk

ABSTRACT

Objective To explore the experiences of accessing and receiving healthcare related to female genital mutilation/cutting (FGM/C) across the life course from the perspective of women and girls who have undergone FGM/C.

Design A systematic review of qualitative research studies using a thematic synthesis approach.

Methods Inclusion criteria were qualitative studies (including grey literature) of any design, from Organisation for Economic Co-operation and Development (OECD) countries, of any date and any language. Sixteen electronic databases were searched from inception to December 2017, supplemented by reference list searching. Papers were screened, selected and quality-appraised by two reviewers using established tools from the Joanna Briggs Institute. NVivo software was used to extract study characteristics and code study findings. An inductive thematic synthesis approach was undertaken to identify descriptive themes and interpret these into higher order analytical constructs. Confidence in the review findings was assessed using Grading of Recommendations, Assessment, Development and Evaluations-Confidence in Evidence from Reviews of Qualitative Research (GRADE-CERQual).

Results Fifty-seven papers (from 55 distinct studies) from 14 different OECD countries were included (50% published within the last 8 years). One-third of studies focused exclusively on maternity care experiences, whereas others covered a range of foci. No studies reported explicitly on girls' experiences or on experiences of health service-led safeguarding interventions. Only three studies addressed psychological care. The synthesis developed 17 descriptive themes, organised into 5 analytical constructs. These related to communication, access to care, experiences of cultural dissonance/integrity, disempowering care experiences and positive care encounters. The themes illuminate significant challenges to obtaining timely and holistic care (especially for deinfibulation), and highlight different ways in which women may experience care as disrespectful, unsafe and disempowering. Key elements of 'culturally safe care' are identified.

Conclusions This review has highlighted key knowledge gaps, especially around (1) girls'/unmarried women's experiences and (2) the impact of recent safeguarding

Strengths and limitations of this study

- This is the first review that goes beyond a maternity care focus to examine women's/girls' experiences of female genital mutilation/cutting (FGM/C)-related healthcare across the life course, enabling new insights into care-seeking and care access, especially around deinfibulation.
- This is an exceptionally comprehensive review due to its wide focus and its inclusion of grey literature and papers in any language.
- The review has used a theoretical model of 'cultural safety' to inform its analytical approach and has assessed confidence in its findings using Grading of Recommendations, Assessment, Development and Evaluations-Confidence in Evidence from Reviews of Qualitative Research (GRADE-CERQual).
- The review process has been informed by strong community involvement and input from a multidisciplinary expert advisory group at every stage.
- Methodological limitations of the included studies make it difficult to develop findings that reflect nuances of experience related to ethnic group, nationality, age or type of FGM/C.

interventions. There is an ongoing need for community engagement, service development and staff training.

PROSPERO registration number CRD420150300012015.

INTRODUCTION

Health services in higher income countries are said to be operating in a context of 'super-diversity'.¹⁻⁵ Whereas in previous decades immigration was primarily from a limited number of countries traditionally linked through former colonial ties, contemporary migration involves groups from many diverse countries across the globe. Superdiversity poses significant challenges to host country health services, in terms of adapting

services and providing culturally appropriate and accessible care to very many different migrant groups with widely different migration histories, languages, health needs and social vulnerabilities.

Female genital mutilation (or female genital cutting)—hereafter referred to as FGM/C—is practised in 30 countries across North and sub-Saharan Africa and in parts of the Middle East and Asia.⁶ In an era of superdiverse migration, European and other high-income countries (eg, USA, Canada, Australia and New Zealand) are becoming home to increasing numbers of women and girls who have experienced, or who may be at risk of, FGM/C (within Europe alone, there are thought to be over half a million women and girls who are FGM/C survivors).⁷ These women and girls come from highly diverse regions with different cultural traditions, contexts and beliefs associated with the practice of FGM/C and where different types of FGM/C are practised. For example, in some communities (eg, within Somalia, Somaliland, Sudan, Eritrea and Kenya), type 3 FGM/C is common involving infibulation (narrowing of the vaginal opening through the creation of a covering seal, formed by cutting and repositioning the labia minora, or labia majora, sometimes through stitching, with or without removal of the clitoris). In other communities, however, type 1 (partial or total clitoridectomy) or type 2 (partial or total clitoridectomy and removal of the labia minora, with or without excision of the labia majora) is more common.⁸ However, it should be noted that these typologies represent biomedical categories and that different communities have their own nomenclature. Moreover, the exact type of FGM/C a woman has experienced may depend to some extent on the skills and experience of the ‘cutter’, and women themselves may not always be aware of which type of FGM/C they have experienced.

FGM/C can involve immediate and long-term psychological, sexual, relational and physical health sequelae.⁹ Health problems may be particularly severe for women with type 3 FGM/C, who require a degree of deinfibulation in order to have sexual intercourse and to give birth. WHO defines deinfibulation as ‘*the practice of cutting open the sealed vaginal opening in a woman who has been infibulated, which is often necessary for improving health and well-being as well as to allow intercourse or to facilitate childbirth*’.⁸ Currently, deinfibulation is recommended for women and girls reporting medical or psychosexual symptoms related to type 3 FGM/C or on request (ie, personal choice). In addition, global guidelines specifically recommend that deinfibulation is undertaken to prevent obstetric complications, although the optimal timing for the procedure (antepartum or intrapartum) is unclear.^{9 9–17} There are compelling clinical reasons however for preferring antepartum deinfibulation, including the fact that it can be performed under local anaesthetic in an outpatient setting, thus reducing costs and risks associated with any emergency procedures that may emerge during labour.¹⁸ In addition, in destination countries where staff may be less familiar with type 3

FGM/C, planned antepartum or premarital deinfibulation ensures that it is performed by trained and experienced professionals.¹⁵

In destination countries, it is essential, therefore, that care pathways are developed that are able to support women’s potential psychological and sexual health needs,^{9–11 19 20} as well as supporting decision making around deinfibulation. In addition, health professionals are called on to play a key role in the prevention of FGM/C, through health education with affected communities and identification of girls who may be at risk.^{20–22} However, as a cultural phenomenon associated with women’s sexuality, gender norms and genital area, FGM/C is a sensitive topic that can be hard for women, communities and health professionals to openly discuss.^{23–25} Recent studies of health sector involvement in the management of FGM/C in destination countries show variable availability of specialist services and staff training.^{21 26} Moreover, several systematic reviews indicate that health professionals may lack knowledge, confidence and competence in managing FGM/C.^{27–30} In addition, there are ongoing calls for greater community involvement in the development of appropriate services related to FGM/C.^{31–33}

Aim

This paper reports the findings of a qualitative systematic review that aimed to explore the experiences of accessing and receiving FGM/C-related healthcare across the life course for women and girls who have undergone FGM/C.³⁴

Rationale

Several prior reviews have been undertaken in relation to this topic; however, we felt a new review was warranted for several reasons. First, most existing reviews have focused on women’s maternity care experiences or maternity-focused interventions.^{35–41} However, as indicated above, a key issue for services is the development of joined-up care pathways that are able to offer women a range of holistic services at different time points. Hence, our review adopted a focus on the life course. Second, it is well known that migrants experience a range of barriers to accessing health services,^{42–46} yet ‘access to care’ has not been a focus of prior reviews related to FGM/C. Third, several prior reviews have included studies from all over the world, thereby bringing together widely different cultural and health system contexts.^{37–39} We felt that a review focused just on ‘destination’ countries would provide findings that were more transferable to these specific contexts.^{47 48} Finally, prior reviews have often limited their searches to English-language papers or to published literature only.^{35 36} By contrast, this review has included any language and a wide range of grey literature sources, and thus is able to offer an extremely comprehensive picture.^{49 50}

Theoretical perspective

This review is broadly informed by the theoretical construct of ‘cultural safety’.⁵¹ Cultural safety derives from critical social theory and argues that traditional approaches to ‘cultural awareness’ or ‘cultural competence’ within healthcare fail to adequately take into account power relations which are historically unequal between migrant or indigenous groups and healthcare providers and services.⁵² Within a cultural competence discourse, for example, it is argued that *‘the power to define the norm and the onus for action to understand and know about another culture fall to the health professional/service’* (p200).⁵¹ As such, migrant groups remain represented as the ‘other’ and may be viewed within a deficit model—as a problem that needs to be fixed—rather than as partners whose knowledge and values can contribute equally to a relationship or to service development initiatives. Culturally safe practice, by contrast, is predicated on relationships of mutual trust and respect. It is a transformative and rights-based approach, seeking to uphold the principles of respect, dignity, empowerment, safety and autonomy.⁵³ Cultural safety envisages the healthcare encounter as a negotiated and equal partnership in which trust plays a central role. The healthcare practitioner’s role is to enable patients to say how a service can be adapted and to negotiate an agreed approach—a key aspect of shared decision making.⁵⁴ The converse of cultural safety is ‘cultural risk’⁵⁵ in that culturally unsafe practices are defined as *‘any actions that diminish, demean or disempower the cultural identity and well being of an individual or group’* (p7).⁵⁶ Cultural safety is not just a feature of individual practice however. It is argued that culturally safe practice must be institutionalised through organisational policies and structures so that it becomes part of mainstream healthcare provision rather than being dependent on individual practitioners who may or may not adopt its values and approach.^{57–59} This theoretical perspective is well suited to areas of healthcare involving stigmatised or sensitive topics, such as FGM/C, and where there is a strong need for community engagement in service development and shared decision making in care delivery.

To date, reflexive consideration of the use of theory in qualitative systematic reviews has received minimal attention. Indeed, some authors have referred to the conceptual process of meta-synthesis as a ‘black box’,⁶⁰ and recent research has highlighted the poor reporting of the interpretive work of synthesis.⁶¹ Yet, as noted by Guba and Lincoln,⁶² there is no such thing as ‘theory-free’ research. Some authors report using a ‘framework synthesis’ approach in which the review is designed to test or expand a pre-existing theoretical framework (which is used as the basis of an initial coding framework to develop key themes).⁶³ In contrast to this relatively deductive approach, in the present review, the concept of cultural safety was used in three ways to support an interpretive inductive approach. First, it acted as one of a variety of ‘lenses’ and perspectives (along with other aspects of our identities and backgrounds) through which

we interpreted the findings of the research studies.⁶⁴ As such, it represented what Maxwell^{39 65} has termed an ‘idea context’ or ‘spotlight’ for the review process in which a theory can help to identify themes that might otherwise be overlooked. Second, as the review progressed, we felt that concepts related to cultural safety remained a ‘good fit’ for helping us to understand and position the review findings within the wider literature.⁶⁶ Hence, we drew on some of its ideas to inform and structure the discussion and recommendations. Finally, we were aware that a common criticism of systematic reviews is their failure to be policy-relevant or transferable.⁶⁷ We felt that framing the discussion and recommendations of the review findings in relation to a theoretical framework might help to enhance the potential transferability of the review findings and guide future research.⁶⁶

METHODS

This qualitative systematic review is reported following the enhancing transparency in reporting the synthesis of qualitative research (ENTREQ) guidelines.⁶⁸ The review was registered in PROSPERO⁶⁹ and the methods are documented in a published protocol.³⁴

Search strategy

An exhaustive and sensitive three-step search strategy was designed by an experienced information scientist (JE). First, we searched 11 electronic databases using a combination of index terms and text-based queries. These were searched from inception to a cut-off date of 31 December 2017. An example search strategy for Ovid MEDLINE is provided in online supplementary file 1. Second, we searched for relevant grey literature drawing on five key resources, Google, Google Scholar and suggestions from the project’s expert advisory group (see [table 1](#) for all resources/databases that were searched).^{70–73} Finally, we hand-searched the reference lists of related systematic reviews and of all the included studies. All retrieved data sets were downloaded into group sets within an EndNote library and duplicates were removed.

Screening and selection

The review included studies from any date and any language that met the following inclusion criteria: (1) empirical research, (2) qualitative research of any design/methodology (including qualitative findings from mixed methods studies), (3) undertaken in an Organisation for Economic Co-operation and Development (OECD) country (OECD was used as a proxy for potential and comparable high-income ‘destination’ countries), and (4) needed to explicitly report women’s or girl’s experiences of seeking, accessing or receiving healthcare associated with FGM/C at any point in the life course and in any clinical setting.

Four team members (RT, CE, JM, GMAH) independently assessed all titles and abstracts against the inclusion criteria. Full-text versions of papers deemed

Table 1 List of databases and resources searched

Electronic databases searched	Date of search
1. Ovid multifile search (MEDLINE, EMBASE, PsycINFO)	10 March 2017
2. POPline (via http://www.popline.org/), 1970–present	10 March 2017
3. ProQuest multifile search	10 April 2017
4. Applied Social Sciences Index Abstracts on ProQuest, 1987–current	26 May 2017
5. Ovid MEDLINE 1948– and MEDLINE In-Process & Other Non-Indexed Citations to daily update	26 July 2017 with monthly alert thereafter (cut-off date for included results 31 December 2017)
6. Ovid EMBASE, 1980–2017 week 11	3 August 2017 with monthly alert thereafter (cut-off date for included results 31 December 2017)
7. CINAHL Plus with Full Text/EBSCOhost to 2017	11 August 2017 with monthly alert thereafter (cut-off date for included results 31 December 2017)
8. Ovid PsycINFO, 1972–March 2017 week 3	14 August 2017 with monthly alert thereafter (cut-off date for included results 31 December 2017)
9. MIDIRS on Ovid, 1971–April 2017	18 August 2017
10. HMIC on Ovid, 1979 to date	18 August 2017
11. Thomson Reuters Web of Science, 1900–2017	18 August 2017
Grey literature sources	
1. British Library Ethos (ethos.bl.uk)	
2. Networked Digital Library of Theses and Dissertations (www.ndltd.org)	
3. National Institute for Health and Care Excellence (https://www.nice.org.uk)	
4. Trove - National Library of Australia (trove.nla.gov.au)	
5. OpenGrey (http://www.opengrey.eu/)	
6. Google	
7. Google Scholar	
8. Experts in the field	

HMIC, Health Management Information Consortium; MIDIRS, Midwives Information & Resource Service.

to be potentially relevant were obtained and assessed. Papers found not to meet the criteria were excluded with reasons noted (see online supplementary file 2). Any areas of ambiguity were discussed with the wider project team. Non-English-language papers found to be relevant on the basis of their English abstract were sent for complete academic translation.

Quality assessment

Study quality was assessed by two reviewers using the Joanna Briggs Institute Qualitative Assessment and Review Instrument (JBI-QARI).^{74 75} Following the guidance of the Cochrane Qualitative Methods and Implementation Group⁷⁶ and the JBI,⁴⁹ reports and theses from the grey literature were appraised in the same way as published peer-reviewed papers.^{70 72 73} Studies were not excluded on the basis of quality, rather the quality assessment was used to judge the relative contribution of each study to the overall synthesis, and to assess the methodological rigour of each study as part of a process of assessing confidence in the review findings.^{77–81}

The JBI-QARI tool was applied to each individual paper and an aggregate score for each was calculated. A criticism of ‘scoring’ qualitative critical appraisals is that it

can be hard to distinguish between the poor conduct of a study and poor reporting, especially where journal word limits constrain the level of detail that can be reported.⁸² In addition, there is no consensus regarding the relative importance of any one domain within an assessment tool over another, and hence whether they should all be given an equal weight or not. In view of these concerns, we chose to adopt a ‘weighting system’ used in previous studies by Higginbottom *et al.*,^{83 84} in which papers were grouped into one of three ‘bands’—high, medium or low—to enable a broad brush evaluation to be made of their relative quality (see table 2).

As an additional strategy for overcoming the potential limitations of solely relying on a checklist to assess quality, we also assessed the ‘richness’ of the studies. This is an approach outlined by Popay *et al.*,⁸⁵ and subsequently operationalised in Noyes and Popay⁸⁶ and Higginbottom *et al.*^{83 84} This approach defines study ‘richness’ as ‘the extent to which the study findings provide explanatory insights that are transferable to other settings’ (p230).⁸⁶ ‘Thick’ papers create or draw on theory to provide indepth explanatory insights that can potentially be transferable to other contexts. By contrast, ‘thin’ papers provide limited or

Table 2 Quality evaluation bands

Quality evaluation	JBI-QARI tool aggregate score	Definition
High	Over 7	A study with a rigorous and robust scientific approach which meets most JBI benchmarks (<i>perhaps 7 or more 'Yes'</i>).
Medium	Between 5 and 7	A study with some flaws but not seriously undermining the quality and scientific value of the research conducted (<i>perhaps 5–7 'Yes'</i>).
Low	Under 5	A study with flaws and poor scientific value (<i>perhaps below 5 of the benchmarks met</i>).

JBI-QARI, Joanna Briggs Institute Qualitative Assessment and Review Instrument.

superficial description and offer little opportunity for generalising. Each paper was assessed against the criteria as set out in Higginbottom *et al* (p5)⁸³ (see [table 3](#)) and categorised as either ‘thick’ or ‘thin’.

Data extraction and assessment of relevance

Study characteristics were extracted by one author (RT) using a modified JBI template and double-checked by CE. During this process, papers were categorised in terms of their relevance to the review question. This assessment was made in order to gain a better understanding of the nature of the body of evidence, and also to facilitate the coding process, as described further below. Study relevance was defined as high, medium or low, as set out in [table 4](#). PDFs of all the included papers were imported into NVivo software, and the ‘findings/results’ and ‘discussion’ sections were coded and analysed.⁸⁷

Data analysis and synthesis

The review adopted a thematic synthesis approach as outlined by Thomas and Harden⁸⁸ involving four iterative stages: (1) indepth reading of the whole papers, (2) inductive line-by-line coding of the findings, (3) grouping the codes on the basis of shared characteristics and meanings into descriptive themes, and (4) interpreting higher order analytical themes.

Studies categorised as ‘thick’, high-quality and ‘highly relevant’ were used as index papers to develop an initial set of ‘open codes’.^{89–101} Two reviewers (RT, CE) undertook this task independently. The codes were then reviewed and compared. Where reviewers had identified and coded the same issue, a code for that issue was agreed for use in coding subsequent papers. A standardised name was identified for the code and a description was

produced. Where reviewers had applied slightly different codes to a concept indicating a different interpretation of meaning, they discussed it and either agreed a shared code (based on a shared understanding of meaning) or created two different codes to reflect two different meanings. The initial set of codes were ‘free nodes’. These were compared, analysed and discussed by the whole team. Where meanings appeared to relate to a similar concept, these were grouped into broad descriptive themes and subthemes. This process created a codebook which was applied to the remaining papers and expanded, refined and modified as appropriate (by identifying new codes and new themes, or by merging and renaming existing codes and themes). The codebook ensured that definitions of codes and themes were explicit and could be easily shared across the team and discussed. More details on this process are given in online supplementary file 3, using the analytical theme of ‘Communication is Key’ (see below) as an exemplar. Analytical themes were evolved through an indepth process of comparing and contrasting the meanings of the descriptive themes, analysing these in relation to how they were, or were not, able to illuminate the review questions, and inferring broader phenomena, categories of meaning or social processes that they related to.^{102 103}

Patient and public involvement

Public involvement was an integral part of the review process and was achieved in three ways. First, the review was conceived and co-constructed from an ongoing partnership between an academic team (CE, RT, GMAH, JM, JE), clinical experts (JA) and a community organisation working on FGM/C-related issues (VN). By undertaking

Table 3 Assessment of study richness

Richness	Operational definition
Thick papers	<ul style="list-style-type: none"> ▶ Offer greater explanatory insights into the outcome of interest. ▶ Provide a clear account of the process by which the findings were produced—including the sample, its selection and its size, with any limitations or bias noted—along with clear methods of analysis. ▶ Present a developed and plausible interpretation of the analysis based on the data presented.
Thin papers	<ul style="list-style-type: none"> ▶ Offer only limited insights. ▶ Lack a clear account of the process by which the findings were produced. ▶ Present an underdeveloped and weak interpretation of the analysis based on the data presented.

Table 4 Assessment of study relevance

Study relevance	Definition
High (specific)	FGM/C-specific healthcare (eg, the study is focused on a specific aspect of care related directly to FGM/C, eg, deinfibulation, childbirth for women who have had FGM/C, psychological care).
Medium (direct)	Other healthcare context (eg, where the study focus is on the maternity care experience of a particular group more generally and where some of the findings relate to the experience of FGM/C).
Low (indirect)	Where the study focus is on general attitudes towards FGM/C and/or experiences and consequences of FGM—and where some FGM/C-related healthcare issues are reported, but are not the main focus of the paper.

FGM/C, female genital mutilation/cutting.

the review we hoped to illuminate women's voices and experiences in order to inform the provision of FGM/C-related healthcare. Second, an expert advisory group was established from the outset, which included FGM/C survivors, activists and health professionals. This group helped with identifying relevant literature and participated in project meetings in which the synthesis and recommendations were formulated. Finally, the review findings were shared with a wide range of FGM/C activist and survivor groups and community organisations, and their feedback helped to shape the final report and recommendations.

Rigour within the analytical process

To ensure rigour of the analytical process, the team sought to identify and understand possible 'disconfirming' cases that might challenge emerging interpretations,¹⁰⁴ and to explore possible subgroup or contextual differences. These processes were aided by the creation of a theme matrix (see online supplementary file 4), in which each theme was mapped to its constituent studies. This helped the team to see clearly how common the theme was among the studies, what kind of study contexts or samples the theme related to, and to explore why it may have been present in some studies but not in others. In addition, as described above, the whole team and wider project advisory group contributed to the evolution of the synthesis, through reading of key papers and providing feedback on the emerging interpretations.

Assessment of confidence in the review findings (CERQual)

Assessment of confidence in the findings of the review was undertaken using the Confidence in Evidence from Reviews of Qualitative Research (CERQual) approach.^{76 77 105–111} Akin to Grading of Recommendations, Assessment, Development and Evaluations (GRADE), CERQual uses assessments of 'concerns' within four distinct domains (methodological limitations, relevance, coherence and adequacy of data) applied to each individual review finding. The CERQual assessment was applied to the descriptive rather than analytical themes, as the latter comprise an aggregation of descriptive themes as well as descriptions of patterns or explanations that are inferred as part of the interpretative process. There is, as yet, relatively little guidance for,

or experience with, applying CERQual to higher level analytical themes or theories.¹⁰⁸

RESULTS

Search outcome

The search outcomes are reported in detail in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow chart in figure 1. Fifty-seven papers met the inclusion criteria, representing 55 distinct studies. Fifty-three papers were in the English language, two were published in German^{112 113} and two in Spanish.^{114 115}

Study characteristics

Due to the large numbers of studies in this review, a highly summarised description of key study characteristics and key methodological assessments is found in table 5. Full details are provided in online supplementary files 5 and 6.

The studies represented 14 different countries, including Australia,^{100 101 116 117} Austria,¹¹³ Canada,^{118–123} Finland,¹²⁴ France,⁹⁰ Germany,^{112 125} the Netherlands,^{126 127} New Zealand,^{116 128} Norway,^{99 129} Spain,^{114 115} Sweden,^{23 94 130–132} Switzerland,^{98 133} Scotland,^{89 134 135} England^{25 92 93 95–97 136–144} and USA.^{91 145–153} The studies were a mixture of older and more recent research, with publication dates ranging from 1985 to 2017. However, 29 papers had been published since 2011; hence, half of the papers reflected a more contemporary context. Over half of the included papers (n=33) were peer-reviewed journal articles.^{23 25 90 92 94 95 97 99 114 116 117 119 122 124 126 127 129–133 136 137 139 144–147 149–153}

Other papers included 13 theses,^{89 91 93 98 112 113 115 118 120 121 123 128 148} 10 unpublished research reports^{96 100 101 125 134 135 140–143} and 1 book chapter.¹³⁸

A large proportion of studies (n=18) focused specifically on women's birth/maternity care experiences.^{89–91 94 95 97 99 117 119 128 130–132 139 145 146 150} Other studies focused on general views on healthcare,^{25 100 101 112 113 115 122 138 140 152 153} general attitudes towards FGM/C,^{23 96 121 123 125 134–136 142 143 151} experiences of sexual/reproductive health services,^{98 114 116 124 133 148} cervical screening,^{126 137 147 149} psychological issues,^{92 93 127} deinfibulation,^{129 144} general practitioner (GP) services,¹⁴¹ identity¹¹⁸ and pain/embodiment.¹²⁰ There were no studies that examined women's




Figure 1 PRISMA flow diagram. FGM/C, female genital mutilation/cutting; OECD, Organisation for Economic Co-operation and Development; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

experiences of surgical reconstruction following FGM/C and no studies that examined experiences of health service-led safeguarding interventions.

The studies were primarily of adult women. Only one study included girls under the age of 18 years but did not report health-related experiences of this group.²³ There were no

studies that focused specifically on healthcare issues related to FGM/C in older (postchildbearing) women. Hence, although the aim of the review was to explore FGM/C across the life course, the studies all focused on generic issues relating to adult women and no further age-specific differentiation could be made. The majority of studies were

Table 5 Summary study characteristics and methodological assessments

Number	Authors (year)	Country	Study focus	Eligible participants and country of origin	Quality assessment	Richness	Relevance
1	Abdi (2012) ¹³⁶	UK	FGM/C and identity for Somali women.	16 Somali women	High	Thin	Low
2	Abdullahi <i>et al</i> (2009) ¹³⁷	UK	Cervical screening.	50 Somali women	Medium	Thin	Medium
3	Ahliberg <i>et al</i> (2004) ²³	Sweden	FGM/C tradition in destination countries.	50 Somali women	High	Thin	Low
4	Ahmed (2005) ¹³⁸	UK	Cross-cultural psychology and women's experience of FGM/C.	56 women (from Somalia and Eritrea)	Low	Thin	Low
5	Ameresekere <i>et al</i> (2011) ¹⁴⁵	USA	Maternity care experiences.	23 Somali women	Medium	Thin	Medium
6	Asefaw (2007) ¹¹²	Eritrea and Germany	Postmigration attitudes towards FGM/C.	31 Eritrean women	Medium	Thin	Low
7	Balden (2013) ⁸⁹	Scotland	Maternity care experiences.	7 women (from Gambia, Ghana and Sudan)	High	Thick	High
8	Ballesteros <i>et al</i> (2014) ¹¹⁴	Spain	Health impacts of FGM/C.	9 Senegalese women	Medium	Thin	Medium
9	Behrendt (2011) ¹²⁵	Germany	Views on FGM/C.	91 women (from Ghana, Nigeria and Togo)	Medium	Thin	Low
10	Beine <i>et al</i> (1995) ¹⁴⁶	USA	Maternity care experiences.	14 Somali women	Medium	Thin	Medium
11	Berggren <i>et al</i> (2006) ¹³⁰	Sweden	Maternity care experiences.	21 women (from Eritrea, Somalia and Sudan)	Medium	Thick	High
12	Betts (2011) ¹¹⁸	Canada	Identity, gender, sexuality and healthcare experiences.	13 Somali women	Medium	Thick	High
13	Bravo Pastor del Mar (2014) ¹¹⁵	Spain	Views on FGM/C.	24 women (from Senegal, Nigeria and Gambia)	High	Thin	Low
14	Bulman and McCourt (2002) ¹³⁹	UK	Maternity care experiences.	12 Somali women	Medium	Thin	Medium
15	Carroll <i>et al</i> (2007) ¹⁴⁷	USA	Healthcare experiences.	34 Somali women	High	Thick	Medium
16	Chalmers and Omer-Hashi (2002) ¹¹⁹	Canada	Maternity care experiences.	415 Somali women	Medium	Thick	Medium
17	Degni <i>et al</i> (2014) ²⁴	Finland	Reproductive health and maternity care experiences.	70 Somali women	High	Thin	Medium
18	d'Entremont <i>et al</i> (2014) ⁹⁰	France	Childbirth experiences.	4 women (from different countries)	High	Thick	High
19	Essén <i>et al</i> (2000) ¹³¹	Sweden	Maternity care experiences.	15 Somali women	Medium	Thin	Low
20	Fawcett (2014) ⁹¹	USA	Views on FGM/C and maternity care.	30 Somali women	High	Thick	High
21	Gali (1997) ¹⁴⁸	USA	Reproductive healthcare experiences.	26 women (from Sudan, Eritrea, Ethiopia and Egypt)	Medium	Thin	Low
22	Ghebre <i>et al</i> (2015) ¹⁴⁹	USA	Views on cervical screening.	23 Somali women	High	Thick	Medium
23	Glazer (2012) ¹²⁰	Canada	Pain and embodiment associated with FGM/C.	14 Somali women	High	Thin	Low
24	Glover <i>et al</i> (2017) ⁹²	UK	Psychological impact, resilience and experiences of FGM/C.	20 women (from Somalia, Kenya, South Africa and Gambia)	High	Thick	High
25	Guerin <i>et al</i> (2006) ¹¹⁶	Australia and New Zealand	Reproductive healthcare experiences.	10 women (from Somalia, Ethiopia, Eritrea, Sudan and Nigeria)	Low	Thin	Low
26	Hill <i>et al</i> (2012) ¹⁵⁰	USA	Maternity care experiences.	18 Somali women	Medium	Thick	Low
27	Hussein (2010) ¹⁴⁰	UK	Views on FGM/C and related services.	8 women (from Somalia and Sudan)	Medium	Thin	Low

Continued

Table 5 Continued

Number	Authors (year)	Country	Study focus	Eligible participants and country of origin	Quality assessment	Richness	Relevance
28	Hussen (2014) ¹²⁸	New Zealand	Views on FGM/C and healthcare experiences.	20 women (from Eritrea, Ethiopia, Somalia and Sudan)	Medium	Thick	Medium
29	Johansen (2017) ¹²⁹	Norway	Experiences and perceptions of premarital deinfibulation.	23 women (from Somalia and Sudan)	High	Thick	Medium
30	Jones (2010) ⁹³	UK	Views and experiences of FGM/C.	6 women (from Sudan and Somalia)	High	Thick	High
31	Khaja (2004) ¹²¹	Canada and USA	Views and experiences of FGM/C.	17 Somali women	High	Thin	Low
32	Khaja <i>et al</i> (2010) ¹²²	Canada and USA	Views and experiences of FGM/C.	17 Somali women	Low	Thin	Low
33	Lundberg and Gerezgijher (2008) ⁹⁴	Sweden	Maternity care experiences.	15 Eritrean women	High	Thick	High
34	Maier (2003) ¹¹³	Austria	Views, experiences and impacts of FGM/C.	3 women (from different countries)	Low	Thin	Low
35	Maternity Action (2014) ¹⁴¹	UK	Maternity care experiences.	23 women (from several countries)	Low	Thin	Medium
36	McNeely and Christie-de Jong (2016) ¹⁵¹	USA	Views and experiences of FGM/C.	13 Somali women	High	Thin	Low
37	Moxey and Jones (2016) ⁹⁵	UK	Maternity care experiences.	10 Somali women	High	Thick	High
38	Murray <i>et al</i> (2010) ¹¹⁷	Australia	Birth experiences.	10 women (from Sudan Liberia, Ethiopia and Somalia)	High	Thick	Medium
39	Norman <i>et al</i> (2016) ¹⁴²	UK	Views and experiences of FGM/C and community engagement.	33 women (from various countries)	Medium	Thin	Low
40	Norman <i>et al</i> (2009) ¹⁴³	UK	Views and experiences of FGM/C, community engagement and services.	40 women (from Sudan, Eritrea and Somalia)	Medium	Thin	Low
41	O'Brien <i>et al</i> (2017) ¹³⁴	Scotland	Views and experiences of FGM/C, community engagement and services.	48 women (from Sudan, Nigeria and Zimbabwe)	High	Thin	Low
42	O'Brien <i>et al</i> (2016) ¹³⁵	Scotland	Views and experiences of FGM/C, community engagement and services.	39 women (from 17 African countries)	High	Thin	Low
43	Palfreyman <i>et al</i> (2011) ⁹⁶	UK	Views and experiences of FGM/C, community engagement and services.	105 women (from Somali, Eritrea, Sudan, Sierra Leone and Gambia)	High	Thick	High
44	Recchia and McGarry (2017) ²⁵	UK	Views on FGM/C and healthcare experiences.	6 women (from Kenya, Nigeria and Ethiopia)	Medium	Thick	Low
45	Safari (2013) ¹⁴⁴	UK	Women's experience of deinfibulation.	9 women (from Somalia and Eritrea)	High	Thin	Low
46	Salad <i>et al</i> (2015) ¹²⁶	The Netherlands	Cervical screening.	46 Somali women	High	Thick	Medium
47	Shaw (1985) ¹⁵²	USA	Views on FGM/C and healthcare experiences.	12 women (from Sudan, Egypt and Somalia)	Low	Thin	Low
48	Shermarke (1996) ¹²³	Canada	Views on FGM/C.	8 Somali women	High	Thin	Medium
49	Straus <i>et al</i> (2009) ⁹⁷	UK	Maternity care experiences.	8 Somali women	High	Thick	High
50	Thierfelder (2003) ⁹⁸	Switzerland	Gynaecological and maternity care experiences.	29 women (from Somalia and Eritrea)	High	Thick	High
51	Thierfelder <i>et al</i> (2005) ¹³³	Switzerland	Gynaecological and maternity care experiences.	29 women (from Somalia and Eritrea)	Medium	Thin	Medium

Continued

Table 5 Continued

Number	Authors (year)	Country	Study focus	Eligible participants and country of origin	Quality assessment	Richness	Relevance
52	Upvall <i>et al</i> (2009) ¹⁵³	USA	Health experiences.	23 Somali women	Medium	Thick	High
53	Vangen <i>et al</i> (2004) ⁹⁹	Norway	Maternity care experiences.	23 Somali women	High	Thick	High
54	Vaughan <i>et al</i> (2014) ¹⁰⁰	Australia	Views and experiences of FGM/C, community engagement and services.	87 women (from Eritrea, Ethiopia, Somali and Sudan)	High	Thick	High
55	Vaughan <i>et al</i> (2014) ¹⁰¹	Australia	Views and experiences of FGM/C, community engagement and services.	50 women (from Togo, Sudan and Kenya)	High	Thick	High
56	Vloeberghs <i>et al</i> (2012) ¹²⁷	The Netherlands	Psychosocial and health experiences.	66 women (from Somalia, Sudan, Eritrea, Ethiopia and Sierra Leone)	High	Thin	Medium
57	Wiklund <i>et al</i> (2000) ¹³²	Sweden	Maternity care experiences.	9 Somali women	Medium	Thin	Medium

FGM/C, female genital mutilation/cutting.

with women from FGM/C practising countries in sub-Saharan Africa, and the majority of these included women specifically from countries in the Horn of Africa, where type 3 FGM/C is most commonly practised. It is likely therefore that the findings included in the review reflect issues that may be specific to these population groups and to the experience of type 3 FGM/C. Only three studies explicitly reported including women from Egypt/Middle East but did not differentiate these women's experience from the rest of the sample.^{138 148 152}

Methodological quality

Full details of assessments of methodological quality, richness (thick/thin) and relevance are provided in online supplementary file 6 (see table 5 for a summary). Thirty papers were assessed as being of high quality,^{23 89–101 115 117 120 121 123 124 126 127 129 134–136 144 147 149 151} 21 as medium quality^{25 112 114 118 119 125 128 130–133 137 139 140 142 143 145 146 148 150 153} and 6 as low quality.^{113 116 122 138 141 152}

A methodological weakness across many studies was lack of discussion of the underpinning philosophical standpoint (question 1 of the QARI tool), making it difficult to assess the congruency of the chosen methodology. Likewise, many studies did not describe any clear methodology (simply stating that they adopted a generic 'qualitative approach'). Such studies tended to have 'thin' or mainly descriptive findings, and it was difficult to judge the congruence of the methodology with the research question and the methods (questions 2 and 3 of the QARI tool). Finally, a limitation across many studies was lack of discussion of reflexivity (questions 6 and 7 of the QARI tool). Given the sensitive nature of FGM/C (and sexuality or migrant healthcare) as a topic, the failure to explore the researcher's own theoretical standpoint or their role, professional background, ethnicity, experience of FGM/C or relationship to the participants is a significant weakness, making it hard to judge the dependability of the findings.¹⁵⁴

Thematic synthesis findings

The findings from 57 papers were synthesised into five analytical themes which represent a synthesis and interpretative analysis of 17 descriptive themes. Table 6 shows their inter-relationship and provides one or two quotes to illustrate each descriptive theme. Hence, the description of themes below does not include quotes. Rather, each analytical theme is explained, followed by an elaboration of its constituent descriptive themes. Due to the large number of studies that contributed findings to each descriptive theme, rather than 'crowd' the text with multiple repetitive references to these individual studies, the reader is referred to a theme matrix (see online supplementary file 4) that shows which studies have contributed findings to each theme.

Analytical theme 1: communication is key

The synthesis revealed that communication was a key interpersonal process that underpinned women's ability

Table 6 Themes, quotations and CERQual assessment

Theme number	Theme heading	Studies (n)	CERQual assessment	Indicative quotes
Analytical theme 1: communication is key				
1.1	Language barriers and interpretation challenges	n=31 23 90 91 95-97 99-101 117-119 124 126-128 130 132-135 137 139-141 143 147-150 153	High confidence	"I wish I had a way to communicate with the doctor directly. There are things that sometimes I want to say or ask but I feel embarrassed saying it through a translator, especially on the phone...They never translate what I'm saying. They tell me that is a shameful question..." (Sweden, p364) ¹⁵³
1.2	'Can't talk, not asked': double silence and cultural taboo	n=37 25 89 91 95-96 99-101 112 115 117 118 120 121 123-128 130 132-135 138	High confidence	"If they had asked me I would have told them, but I was never asked...I was worried and scared during my first pregnancy because I have heard people talk about the pain but I didn't know how to tell them...I was shy." (Scotland, p23-24) ⁸⁹
1.3	Cultural (n)sensitivity	n=34 25 89 91 92 96-101 112 115-122 125 127 128 130 132 135 136 138-143 146-148 153	Moderate confidence	"The doctor was not comfortable with me. He asked if I enjoy violent sex with my husband. I don't feel comfortable to talk about sex even with females—yet here is a male doctor examining my genitals and making mean comments." (USA, p274) ¹¹⁹
Analytical theme 2: access to care: influenced by an interaction of multilevel community and health service processes				
2.1	Influence of cultural norms	n=35 93-95 97 99-101 112 113 117-121 124 126-130 132 133 137 138-141 143-147 149 150 153	Moderate confidence	"Even though we had FGM type 2 or 3, had a small vaginal opening and were virgins, we did not ask for deinfibulations before marriage because that is shameful in our culture. Most of us who have FGM had problems with penetration following marriage." (New Zealand, p43) ¹²⁸ "In my country normally...people don't go...usually people go to a psychologist if it is something like erm... maybe chronic issues that they cannot solve at all...or sometimes if it is abnormal... They think if you go to a psychologist you are mad or crazy." (England, p6) ¹³³
2.2	Influence of the family	n=19 90 91 93 94 97 99 100 113 118 121 124 126 130 132 136 144 149 150 153	Moderate confidence	"I know my friend she wants to do that (deinfibulation) but her husband doesn't let her...I told her, this is good for you - go and do. She said ok but her husband said no...Maybe if she did this (deinfibulation), it would have been a big problem in the family." (England, p157) ¹⁴⁴
2.3	Knowledge and information about FGM/C services	n=34 92 93 95-97 99-101 113-119 121 126 128 130-135 137 139 142 143 145 148-150 152 153	Moderate confidence	"They need to have someone that talks to you before and while you have a baby. I would have wanted someone to help me and tell me what was going to happen but I never got that... You know someone who was specially trained so you only see people who understand FGM." (England, p233) ⁸²
2.4	'Hit and miss' care	n=38 89 91 92 94-97 99-101 112 113 115 117-120 123-125 127 128 130-135 139-143 147 148 150-152	High confidence	"Sometimes people go to one doctor after another describing their problems to them. They do not understand so they cannot find a solution. There was a woman who was suffering from bleeding and pain. When she went to the GP, he told her it was normal. He did not know how to treat her. She kept going several times and at last they opened her." (England, p41) ¹⁴³
Analytical theme 3: cultural and bodily dissonance: striving for cultural and bodily integrity				
3.1	Moving from normal to different	n=40 89 91-94 96 97 99-101 112 113 115-120 122-130 132-135 140 141 143 144 148 150-153	High confidence	"Okay, now I'm missing something that other women has... Well the first shock was to find out that people here are not circumcised... You have to deal with that...I'm not the same as women here...and then I started to kind of - you grieve a bit, that 'what is this thing that has been taken away from me?'...For me it has caused me lots of problems because when I went to health services, the minute that I say 'I was circumcised' it just create huge barrier between them and me, like you know - suddenly I'm from another planet." (Australia, p23) ¹⁰⁰
3.2	Threat to the self: reliving FGM/C pain during clinical interventions	n=28 89 92-94 95 99-101 112-114 119 120 126-128 130 133 134 136 137 140 143 146 149 151 152	Moderate confidence	"You always remember it. They decide to cut me, and when she went to do it she didn't explain to me and I panic. I nearly kick her. Two of the nurses held me down. That was it...I was a little girl again being held down." (England, p228) ⁹²
3.3	Being opened: complexities around deinfibulation	n=28 89 92-97 100 101 112 113 115 116 118 120 121 124 128-130 133 134 143 144 146 148 150 152	Moderate confidence	"She [Somali lady] hadn't talked to friends about it, just the midwife...She said: 'When I met the midwife she told me a lot about FGM. The midwife said to me 'if you want I can open it for you. If you don't want that I won't'. We talked about it a lot. After four months I tell her 'I want you to open it for me. Do me a small operation.'" (Scotland, p31) ³⁴

Continued

Table 6 Continued

Theme number	Theme heading	Studies (n)	CERQual assessment	Indicative quotes
3.4	Being changed: complexities around reinfibulation	n=19 92 94 96 98 100 101 112 113 115 116 118 124 129 130 134 144 146 148 152	Moderate confidence	"I became a victim in Sudan already when I was 4 years old: I had no choice. Now I have to become a victim again after delivery, when the midwives refuse to resuture me. I just ask for a few stitches, not to have an open wound." (Sweden, p53) ¹³⁰
Analytical theme 4: disempowering care encounters				
4.1	Being exposed and humiliated	n=40 25 89-92 95-97 99-101 112 113 115-120 122 125-128 130 132-138 140-143 146 148 151 153	High confidence	"Me - I was installed with my legs spread apart - they all came to see...and I didn't see any reason why they came to see me except for the fact that I was excised. All those people who came, who entered into my intimacy without my authorization, who didn't ask me if bothered me...I felt that it was an aggression." (France, p308) ⁸⁰ "They showed no respect to me, I understand now why people don't want to talk about it to the midwife. I will tell anyone not to tell them about being circumcised." (Scotland, p23) ⁸⁹
4.2	Being judged and stereotyped	n=36 23 25 89-93 96 97 99-101 112 113 115-119 122-125 127 130 132 133 139-143 145 147 148 153	High confidence	"One participant...described that a German gynaecologist (a woman) looked at her with so much pity and irritation after the physical examination that she felt for the first time in her life that she was not normal...she felt that she was treated like a victim who could not be talked to rationally. The participant decided...to abstain from consulting German doctors." (Germany, p83) ¹²⁵
4.3	Lacking choice, power and control	n=32 89-92 94 95 97 99-101 112 113 116-121 123 125 127 130-134 139 141 143 145 146 148 150	High confidence	"I can't say that the birth went well. The midwife said that I couldn't give birth without being cut. otherwise the baby won't come without a caesarean section or I can lose the baby. Nobody asked me what I thought. I was there in pain...without pain relief, nothing." (France, p308) ⁸⁰
4.4	Feeling unsafe and vulnerable	n=46 23 25 89-97 99-101 112-121 123 125 128 129 130-135 139-141 143 145-148 150-153	High confidence	"They didn't know what they were doing, and uh, they, you know they had to give me more cuts then in an awkward way...I was telling the doctor to cut the FGM and totally he ignored me. And he was cutting in the sides instead of cutting the front. It was very degrading and I am very distressed to think about it." (Australia, p25) ¹⁰⁰
Analytical theme 5: positive care encounters				
5.1	Trusting and appreciating providers and the system	n=34 90 92-96 99-101 112-115 117-120 123-125 127 128 130-132 134 135 139 143 145 146 148 150 153	High confidence	"I was lucky when I met a midwife in Sweden who knew about circumcised women. This was a great help to make me feel secure because it was my first time to be pregnant and to live far from my parents and family." (Sweden, p219) ⁸⁴ "I know a friend that had to be opened before her delivery. She was lucky because she had found a midwife who knew about FGM - and she booked her to have the operation before the delivery." (England, p15) ¹⁴⁰
5.2	Voicing healthcare needs and preferences	n=45 25 89-97 100 101 112 113 115 116 118-120 122-124 127-130 132-135 137-143 145-148 150-153	High confidence	"A while ago, all the women used to choose caesarean to give birth but now the health services know about women who are circumcised. They know of their situations and the complications they can have during maternity like bleeding and that they need help with the opening before the baby comes. When the health services don't know about your condition, they refer people to other services. There are some who are understanding, especially if different communities come to attend the clinics or hospitals, as they get more oriented on the different cultures of the communities." (England, p43) ¹⁴³

CERQual, Confidence in Evidence from Reviews of Qualitative Research; FGM/C, female genital mutilation/cutting.

to seek care, to obtain care and to have a positive care experience. Conversely, difficulties in communication negatively impacted all of these dimensions. Communication about FGM/C was characterised by issues that might be expected within migrant women's encounters with new host country healthcare systems—such as language barriers within consultations. However, communication about FGM/C presented an additional layer of complexity related to the fact that FGM/C was seen by both women and healthcare providers as a deeply personal, private and sensitive issue touching on a range of taboos associated with gender, culture and sexuality—all issues that in many societies are shrouded in secrecy and silence. When women encountered providers who were able to overcome taboos and cultural differences, it forged bonds of trust and facilitated an environment in which women were able to talk about their FGM/C and to explore options for care. However, the majority of studies reported challenges and problems with communication and interpretation which led to women being unable to talk about their problems or to explore appropriate treatment options, led them to avoid healthcare in general and generated a sense of mistrust in the system. This theme mainly related to maternity care interactions, but also to communication about cervical cancer screening.^{100 101 126 134 141}

Theme 1.1: language barriers and interpretation challenges

The majority of studies (n=31) reported that for women who did not speak the host country language well, communication about FGM/C was hindered by language barriers and problems with accessing interpretation support that was appropriate for discussing highly sensitive and personal issues. Language barriers meant that women were unable to form a trusting relationship with their provider, express their needs adequately or understand information or advice. This led to frustration and increased anxiety, especially in a context where consultation times were often limited.^{128 139} In some cases, women felt that they had had poor clinical experiences as a direct result of their inability to communicate about FGM/C.^{23 128 134} In other examples, women avoided mentioning their FGM/C at all because they knew they lacked the language to explain.^{128 133 140}

Theme 1.2: 'Can't talk, not asked': double silence and cultural taboo

As a private, sensitive and taboo issue, communication around FGM/C was hindered by a double silence. Women reported that FGM/C was rarely discussed within their own communities, and likewise they found it hard to discuss it with health providers, especially if these (or the interpreters) were male. For this reason, women preferred the topic to be raised by health providers; however, a commonly reported finding was that, even when women appropriately accessed relevant services (eg, GPs or antenatal services^{95 142 143}) and might appreciate the general care offered,^{115 124} healthcare providers often did not ask about FGM/C, even when the consultations were for

pregnancy-related check-ups. The consequence was that FGM/C was sometimes not identified or discussed until women presented in the labour room, so opportunities for care planning or birth planning were missed.⁸⁹ In addition, some studies reported that women found communication to be easier when supported by a knowledgeable community advocate or confident peer.^{89 96 118 141 147} Other studies emphasised that open communication was predicated on being able to develop a trusting relationship with the health professional, which in turn required time and continuity of care.^{95 100 101 139}

Theme 1.3: cultural (in)sensitivity

This theme was reported by 34 studies in which women described experiencing comments and questions from health providers that were perceived as clumsy, insensitive or intrusive. As a result, women reported feeling ashamed, scared or stigmatised. Such encounters inhibited women from talking about their FGM/C, and in some cases led to them avoiding examinations or healthcare services.^{112 148} By contrast, culturally sensitive communication was greatly appreciated and enabled women to openly discuss their FGM/C and related questions.^{90 95 135}

Analytical theme 2: access to care: influenced by an interaction of multilevel community and health service processes

The review found that access to care related to FGM/C was influenced by an interaction of factors operating at individual and community levels, as well as at health system and service levels. At individual and community levels, studies showed that care-seeking was influenced by cultural norms around health and sexuality, collective approaches to decision making, and the level of knowledge and information that individuals and communities had about FGM/C and health service availability. These factors influenced whether or not a woman would seek care, at what time period and where. However, the studies also showed that care-seeking choices were shaped by the types of services available and that outcomes were influenced by the extent to which the health system, health services and health practitioners were ready and able to provide the range of support that women may need at different points in their lives. The review showed that there were challenges within communities in terms of seeking care and finding the right services. However, even when services were appropriately accessed, care was sometimes haphazard and suboptimal due to variable levels of staff expertise and inconsistent and unclear referral pathways, policies and procedures.

Theme 2.1: influence of cultural norms

Women's care-seeking in relation to FGM/C was strongly influenced by wider cultural norms around sexuality and health, including cultural norms on the importance of premarital virginity, avoiding male health providers and lack of familiarity with preventive care-seeking. These factors often led women to avoid seeking care unless

symptomatic or pregnant. Thirty-five studies contributed to this theme.

Women reported that strong cultural imperatives valuing premarital chastity and virginity meant that they would avoid procedures requiring gynaecological examinations (eg, Pap smears^{101 126 143 149}), and especially for those with type 3 FGM/C the general norm was to avoid seeking deinfibulation prior to marriage and pregnancy due to the cultural pressure to maintain chastity, prove their virginity and to remain 'closed' for their husbands.^{95 98 128 129 149 152} This norm was so strong that some studies reported younger women wishing that they could undergo deinfibulation, experiencing unpleasant symptoms and being aware of services, but feeling unable to resist community pressure.^{93 129 144} Some exceptions to this norm were reported, but mainly in the context of needing help for particularly difficult or painful symptoms.⁹⁵

Care-seeking in relation to mental health issues associated with FGM/C was infrequently reported, and a few studies suggested that this may also be linked to cultural norms in which mental health is still often viewed as a stigmatised issue and, like FGM/C itself, is seen as hard to talk about.^{93 123 128 143 150}

Theme 2.2: influence of husbands and the family

Nineteen studies reported that access to care in terms of healthcare decision making, especially among Somali women, was strongly influenced by the views and advice of the wider family and peer group. In the maternity context, this had an impact on decision making around uptake of antenatal care and caesarean section.^{91 93 95 118 132 149 150} The family also influenced decision making around FGM/C specifically. For example, as mentioned above, women considering premarital deinfibulation reported that they might avoid informing their direct family members as they were aware that they would otherwise come under strong pressure to change their mind due to fears over their future marital prospects.^{96 144} For married women with type 3 FGM, studies indicated that the views of husbands were of paramount importance and that women generally felt that they needed their husband's permission to undergo deinfibulation (outside of the context of delivery).¹⁴⁴

Theme 2.3: knowledge and information about FGM/C services

Women's knowledge of, and familiarity with, health services was variable. In some cases, women lacked familiarity with the host country health system, and this impeded their ability to access care in general. In other cases, women reported being quite familiar with maternity and primary care services, but lacked knowledge and information of FGM/C-specific specialist services (especially non-maternity-related services such as counselling) and where and when it may be appropriate to seek help.^{93 134 143} One study from the UK suggested that this

might be a particular issue for women living in low prevalence areas.¹⁴²

Theme 2.4: 'Hit & miss' care

When accessing healthcare, women reported that (1) the identification of FGM/C and (2) provision of appropriate treatment or referrals could be a 'hit and miss' process, depending on individual provider characteristics and practices, rather than being a result of standardised organisational systems and processes. For example, many studies cited situations where opportunities to identify FGM/C during a consultation were missed, which was attributed to providers lacking awareness of FGM (or related specialist services), and hence failing to ask the right questions, or to undertake an examination or to make timely referrals.¹⁴³ Maternity services were more likely to have knowledgeable providers and responsive reporting and follow-up systems, but even here women described situations where they felt their FGM/C had been mismanaged due to lack of awareness and lack of appropriate referrals.^{96 115 141 143} The most common issue was failure to have been asked about FGM/C so that it was not included in birth planning discussions and was identified only once labour had started.⁹² This theme was reported in 38 studies across different countries and time periods.

Analytical theme 3: cultural and bodily dissonance: striving for cultural and bodily integrity

This analytical theme refers to changes in and challenges to women's sense of cultural and bodily identity and integrity as a result of their experiences associated with FGM/C in the host country, and particularly through their encounters with the healthcare system (manifested, for example, during experiences of clinical examinations, childbirth, deinfibulation and reinfibulation). The theme describes women's responses to experiences of dissonance, whereby their decision making and actions can be understood as a desire to maintain a sense of cultural and bodily integrity. This theme highlights differences in, and changes to, cultural meanings and values related to FGM/C and how these differences and changes influence the way that women and health professionals define health issues, make decisions around FGM/C care and experience that care. When mutual interpersonal cultural understanding was not achieved, women perceived suboptimal care and distrusted the health system and their health providers. If women felt understood, they felt safe and reassured, resulting in a sense of bodily and emotional integrity.

Theme 3.1: moving from normal to different

Some women described becoming aware of FGM/C as something that is 'different' only once they moved to another culture. At this point, some started to become aware of the suffering and symptoms that their FGM/C may have caused them. They started to question and to resist previously taken for granted aspects of their culture

and to feel uncomfortable with their own bodies and sexual identity.¹⁰⁰ However women also reported feeling shame and anger at being labelled as ‘different’ and ‘mutilated’ by the dominant discourse in the host country and by health professionals, and felt that their culture was misunderstood.^{90 96 136 137} This experience of dissonance made some women feel reluctant to mention FGM/C to health providers or caused them to feel extremely uncomfortable during healthcare consultations, thereby limiting the opportunity to discuss the issues more openly and to achieve greater mutual understanding.^{91 100 130} This theme was very common—reported in 40 studies.

Theme 3.2: threat to the self: reliving FGM/C pain during clinical interventions

Many studies (n=28) reported women’s experiences of gynaecological clinical interventions and childbirth as events which threatened their sense of bodily and cultural integrity. Women described great pain, suffering, fear and apprehension around clinical interventions—and related this to their previous healthcare experiences, and especially to reliving the original trauma they had experienced during their FGM/C. These emotions were exacerbated in healthcare encounters where women felt a loss of control or lack of respect. Negative experiences could lead to women avoiding attending the hospital until the last minute as a way of avoiding medical interventions.¹⁵⁵

Theme 3.3: being opened: complexities of deinfibulation for women with type 3 FGM

The experience of dissonance and quest for integrity was also seen in views and experiences around deinfibulation and was reported by 28 studies. As noted above, for cultural reasons, ‘being opened’ medically was seen as necessary primarily after marriage and primarily in the context of pregnancy and childbirth, rather than before.^{121 122} There were no studies that explicitly or exclusively focused on the views or experiences of unmarried women, but some studies reported examples of unmarried women where the need to maintain cultural and bodily integrity outweighed the experience of physical pain or discomfort, even when women knew about the option to seek surgical deinfibulation.^{93 128 129} There were few reported exceptions to this rule, although one study suggested that for some women, finding out about the option for surgical deinfibulation was empowering and deinfibulation was subsequently seen as a way of asserting control over their bodies and lives.⁹⁶

Most studies reported that women preferred to be ‘opened’ during labour to avoid the pain and trauma of being cut twice.^{91 95 98 120 129 144} However, some studies suggested that this view could change towards the medically preferred option of antepartum deinfibulation, if there were opportunities for women to have appropriate discussions, conducted in a sensitive way in the context of a trusting relationship with a healthcare provider.^{130 134} Hence, given that FGM/C is often not discussed during healthcare consultations, women’s current reported

preferences around deinfibulation timing may also reflect the fact that they may not be receiving the right information at the right time, and thus may not be in a good position to make informed decisions on this issue.^{96 112 128 134}

Theme 3.4: being changed: complexities around reinfibulation

A consistent finding across the studies within this theme (n=19) was that ‘being opened’ could involve significant emotional, physical, social and relational adjustments as women’s bodies became changed. For some women, deinfibulation was firmly welcomed, although still required a period of ‘getting used to’ a different way of looking and feeling.^{100 144} For other women, however (especially women from Sudan where it is a cultural norm in some regions to be closed again after delivery), there was ambivalence about these changes and some women requested to be ‘closed again’ to varying degrees.^{94 96 100 118 128 129 144 146}

In most OECD countries, reinfibulation is illegal, and some studies reported women feeling very upset when their requests were denied by health professionals, experiencing this as a denial of their agency and integrity.^{112 116} Consequently, women in one study reported seeking medical deinfibulation during holidays back home.¹³⁴

Analytical theme 4: disempowering care encounters

In many situations, the care received by women in relation to their FGM/C was experienced as fundamentally disempowering. Women described experiencing negative attitudes or behaviours from health professionals and poorly managed clinical interventions in which they felt they were not respected or listened to. Many studies reported examples of care encounters that left women feeling retraumatised, voiceless, without power to question interventions and vulnerable in the hands of unprepared system. Such experiences caused women to feel unsafe and disempowered. There were many examples where women felt that care had gone wrong or where experiences had been poor, and that these situations could have been avoided with better communication, cross-cultural understanding and sensitivity. This analytical theme is made of four descriptive themes that cut across all countries, time periods, population groups and care contexts.

Theme 4.1: feeling exposed and humiliated

Across many studies (n=40), women reported that the reactions of healthcare professionals to their FGM/C had left them feeling ashamed, objectified, humiliated and exposed. This was primarily the case with healthcare professionals who were unfamiliar with FGM/C and who were encountering it for the first time. In these situations, women reported that healthcare professionals often reacted with extreme shock when they saw that a woman had been cut. Some tried to hide their shock and avoided mentioning anything at all, making the encounter feel uncomfortable. Others openly expressed horror or

disgust or reacted insensitively by asking intrusive or inappropriate questions. In many cases, other colleagues were called in to be consulted or to ‘have a look’ (sometimes without consent^{119 130}), causing women to feel that their privacy had been violated and that they had been put on display. Such experiences left women feeling angry, distressed or as if they had been reviolated.^{90 91} In some cases, these experiences left women wanting to avoid further contact with services and/or feeling unable to discuss their care any further as it was clear that the provider had limited knowledge about FGM/C.¹⁴⁸ Several studies made the point that negative experiences of a particular service or provider would then be discussed within peer and community networks, and hence could have an impact beyond the individual woman, potentially affecting the way others in the community might engage with care.^{89 91 92 112 122 124 127}

Theme 4.2: feeling judged and stereotyped

In many studies (n=36) women reported feeling judged and discriminated against within some of their healthcare encounters. They reported feeling that health providers made negative assumptions about them and provided suboptimal care based on racial, religious or other stereotypes and misconceptions about their culture in general, as well as about FGM/C. Such experiences caused distress, anger and avoidance of the health provider or service.¹³⁰

Theme 4.3: lack of choice, power and control

In this theme, reported by 32 studies, women described experiences where they felt they had lacked choice or control within the healthcare encounter, especially in the maternity setting and around key interventions such as caesarean sections or episiotomies. Lack of control was experienced in terms of feeling excluded from healthcare decision making, not being informed, not being listened to, feeling at the mercy of ‘the system’ and unable to express their needs. Women’s attempts to assert control were sometimes construed by health professionals as being ‘difficult’.^{91 92 119 145 150} In labour, women described that FGM/C was often not discussed at all or that they were not consulted about interventions such as deinfibulation or caesarean section. This led to suspicion and fear, and was particularly reported in studies conducted with the Somali community.⁹⁰

Theme 4.4: feeling unsafe and vulnerable

Many studies (n=46) reported care encounters where women felt unsafe and vulnerable, primarily when they were treated by providers who they perceived to be inexperienced in dealing with FGM/C, and when they faced language barriers or were unable to express their own preferences. Such encounters often led to perceived poor-quality care or poor clinical outcomes.¹³⁹ The consequences ranged from providers failing to recognise, identify or discuss FGM/C,⁸⁹ to situations where women felt their care had been adversely affected as a direct result of poor provider skills (eg, having unnecessary caesarean

sections or extensive perineal tearing).^{23 25 89 94 95 99 112 115 128 143 153} Women endured such situations feeling unsafe and highly vulnerable, often describing painful and traumatic experiences.¹⁰⁰ Such feelings of vulnerability were heightened for women who lacked social support.^{90 95 130 140} In contrast, women reported feeling safe and highly reassured when they encountered providers who appeared to be knowledgeable or experienced.^{89 92 94 96 100 101 135 139 143}

Analytical theme 5: positive care encounters

As seen above, the synthesis has revealed many reports of poor care and difficult experiences in relation to FGM/C. However, many studies reported a mixture of experiences, including both good and poor care. High-quality care was conceptualised by women as care that is safe, respectful, culturally sensitive and compassionate. The essential process underpinning the experience of ‘good’ care was having a sense of trust—both in the individual provider and also in the ‘system’ as a whole. Trust was essential for, and developed from, positive care experiences. Positive experiences were associated with the development of a good relationship with health professionals who were perceived to be clinically knowledgeable and culturally sensitive, facilitating open communication and a sense of being in safe hands. Trust in the ‘system’ was linked to women’s appreciation of the availability and accessibility of services (compared with their home countries), but particularly when services were perceived to be prepared to be responsive to community needs (eg, provision of appropriate interpretation services) and to manage issues relating to FGM/C (eg, by having specialist services or specialist providers).

Theme 5.1: trusting and appreciating providers and the system

This theme, reported in 34 studies, showed that women appreciated the good medical services available in the host country. Women also expressed great appreciation for providers who made them feel safe and respected. Such providers were described as knowledgeable and experienced, who treated women with respect, who understood their individual needs and who involved them in their care. Women described these characteristics as facilitating trust and leading to open communication about FGM/C. For example, they were more likely to engage in conversations about their care (eg, birth planning), and in some cases were more likely to follow current medical guidelines, for example, on issues such as deinfibulation timing.¹⁴⁰ A few studies explicitly linked the ability to form good relationships to services that enabled continuity of care.^{95 139}

Theme 5.2: voicing healthcare needs and preferences

Women’s recommendations for good-quality, safe FGM/C-related care and services they could trust were reported in 45 studies. They included (1) interpersonal provider characteristics and behaviours (such as providers being willing and able to talk about FGM/C, providers being skilled and knowledgeable around

FGM/C, and providers offering culturally sensitive and respectful care) and (2) service organisation issues (such as having specialist services for FGM/C, being given information and awareness about FGM/C-related services, and involving women and affected communities in FGM service development).

Confidence in the review findings

Confidence in the review findings (as assessed by GRADE-CERQual) ranged from high to medium, and there was strong consistency in the findings across countries, population groups and clinical contexts (see table 6 for the final CERQual assessment for each theme and see online supplementary file 7 for the full details of the CERQual evaluation).

DISCUSSION: ACHIEVING CULTURALLY SAFE CARE FOR WOMEN WITH FGM/C

The review shows that FGM/C—literally and symbolically—embodies complex processes of cultural change and cultural interaction that are occurring as a result of globalisation, migration and superdiversity. These processes manifest themselves very concretely in healthcare in terms of influencing how individuals seek care, how care is delivered and experienced by all involved, and how services are configured.^{2 156 157}

This review has shown that, when obtaining care for FGM/C-related issues in a host country, women face common challenges that affect many groups of migrants, but that these are exacerbated.^{46 84 158–161} Silence, secrecy, stigma and lack of familiarity with FGM/C within the system combine and act as obstacles to identifying FGM/C or to providing women with appropriate care. The care for FGM/C described in this review exemplifies the challenges of achieving ‘cultural safety’ in health services. Many of the experiences reported in the review describe a situation of physical but also cultural ‘vulnerability’ or ‘risk’ whereby women felt physically unsafe but also disrespected in terms of their cultural identity and bodily integrity and uninformed in their care. Many of the themes in this review referred specifically to maternity care contexts, and there are striking similarities with themes identified in studies across the world that highlight disrespect and abuse in maternal healthcare.^{35 36 162} However, as this review shows, the same challenges also appear to exist for women with FGM/C in primary care and other health settings.^{163–165}

This review has also shown that, in some circumstances, women and providers can overcome these obstacles. Key to this are knowledge of services, community engagement, the availability of specialist services, continuity of care, shared decision making, addressing language barriers, and providing care that is person-centred, culturally sensitive and respectful.^{44 159 166 167} Similar recommendations have also been made in other studies exploring the key elements of ‘respectful care’ in a variety of settings.^{168 169}

Figure 2 outlines a model, derived from the review findings, of four key elements of ‘culturally safe’ care for women/girls with FGM/C, set within a wider context of superdiversity.

Recommendations for culturally safe care

As per the model in figure 2, the review recommendations fall into four overlapping areas.

Overcoming barriers: information, awareness and community engagement

The review showed that there is a need for communities to be more aware of FGM/C-related services and of potential interventions. However, it also highlighted the challenges that women may have in discussing FGM/C within their family or community and with a health provider. This finding highlights a need for ongoing community involvement and engagement to raise awareness of services and provide support to women who may wish to access help (especially for unmarried or non-pregnant women).^{31 32} The review suggests that service models which involve community advocates or community liaison workers may be particularly helpful in encouraging women to access care and to overcome communication difficulties with providers. Similar models have been found to be effective in other areas of healthcare.^{170–174}

Breaking the silence: open communication and shared decision making

Given the communication challenges around FGM/C, the review findings strongly suggest that the onus of ‘breaking the silence’ may lie with the healthcare provider, with women stating that they expected and wanted health professionals to raise the issue. It may be appropriate for services to consider routinising questions around FGM/C in key clinical settings, as has been done for other ‘sensitive’ clinical issues such as domestic violence or HIV.^{175–178} In addition, given the particular global concern for women’s (and their babies’) well-being related to timing of deinfibulation surgery (for women with type 3 FGM/C), the review findings strongly suggest that women appreciate, and would benefit from, more discussion, information and advice—at different stages in their life (before and after marriage) and both predeinfibulation and postdeinfibulation.

Provider competence and confidence

The review highlighted that providers may lack the confidence or cultural competence to raise the subject of FGM/C in an appropriate and respectful way, potentially missing opportunities to identify FGM/C in a timely manner but also missing opportunities to discuss prevention/safeguarding.³⁰ Communication challenges can be addressed through staff and interpreter training.¹⁷⁹ Likewise, the review showed that providers would benefit from additional training in associated clinical skills such as deinfibulation. The evidence base for best practice in health worker training around FGM/C is still weak¹⁸⁰; however, as per WHO guidelines,²⁰ it is suggested that




Figure 2 Culturally safe care for female genital mutilation/cutting.

basic training around FGM/C should be embedded in preservice curricula for all health professionals, as well as being offered as continuous professional development for providers in more specialised roles.

Getting the model of care right: appropriate systems, services and care pathways

The review suggests that women sometimes receive suboptimal care due to their FGM/C being missed or because of a lack of appropriate referral and reporting pathways. Some countries have made great strides to develop clear processes in this area (eg, UK), but in others service provision remains ad hoc.²¹ As per WHO guidelines,^{9 20} the review affirms the ongoing need to develop holistic care pathways.^{13 26 181–184} The review findings suggest that women have better experiences in places where there are specialist services with confident and competent staff, and that women in lower prevalence or rural areas may be at a particular disadvantage. Hence, it is important for regions to develop service models that can address these inequalities. In addition, the review strongly suggests that respectful open discussions about FGM/C are more likely to happen in a context of a trusting relationship, and that such relationships are facilitated through service models

that implement continuity of care approaches and shared decision making.^{163–165 185}

The review has highlighted the ongoing need for the availability of mental health assessment and support where required. In particular, the review showed that for some women affected by FGM/C, childbirth and other clinical interventions could be traumatising, triggering flashbacks of their original cutting, creating profound anxiety and distress, and potentially leading to poor care experiences or avoidance of services.^{186 187} The review was unable to illuminate in any further depth how such women might best be identified or treated, or which women might be most at risk. This is an important agenda for future research.^{13 39 188–192} However, most maternity care services have well-established processes in place for dealing with highly anxious women or women who have experienced trauma. It would seem important for services to ensure that these interventions are also available for women affected by FGM/C.^{35 193}

Strengths and limitations

As mentioned above (under the Rationale section), previous reviews have primarily focused on care

experiences within maternity contexts.^{35 36 40 41} Our findings on care experiences resonate strongly with these existing reviews. However, we feel that our review has been able to expand on current knowledge in several ways. First, it adopted an exceptionally comprehensive and exhaustive search, and we are confident that the majority of relevant papers were identified. The inclusion of grey literature as well as studies in other languages significantly broadened the evidence base (eg, the existing reviews on maternity contexts included between 4 and 16 papers, whereas this review included 57).^{35 36} It is our contention that this wider evidence base, as well as the broader review focus on all aspects of the life course, has enabled a more nuanced understanding to emerge around FGM/C-related health experiences. In particular, our review has identified new findings around (1) factors that influence care-seeking and access to care (as well as the care experience itself), (2) decision making around deinfibulation surgery, (3) community perspectives on positive care experiences and suggestions for service improvement, and (4) being able to demonstrate the importance of primary care as well as maternity contexts for discussing and identifying FGM/C. The theme matrix presented in online supplementary file 4 provides a visual means for exploring in more detail how the wider evidence base has contributed to particular review findings, and thus how it has enabled new or more nuanced themes to be identified.

Another strength of the review is its exclusive focus on OECD countries, which ensures that its findings are highly relevant to health service development in most destination countries.

The main limitation of the review is that the findings inevitably reflect the methodological limitations of its included studies. Many papers displayed a homogenising tendency in terms of the affected communities as well as FGM/C types. For example, in papers that had mixed community samples, there was little attempt to explore cultural differences in relation to FGM/C. Likewise, it was not always clear which type of FGM/C was being studied. In research with Somali or Sudanese communities, the assumption can reasonably be made that their focus was FGM/C type 3. However, in studies with mixed samples, there was very little differentiation on the basis of FGM/C type. As a result, there is limited specific knowledge on the needs or experiences of women with other FGM/C types.

Evidence gaps and future research

The review has identified five key gaps in the evidence base around FGM/C, all of which are priority areas for future research. First, there were very few studies that focused specifically on mental health needs. Second, there is a lack of studies on girls' or unmarried/non-pregnant women's experiences around FGM. Important areas for future investigation among this group are care-seeking and decision making (especially

around deinfibulation timing). Third, there is a lack of knowledge on potential health needs or experiences of older women; hence, this is an area that may require further investigation. Fourth, there were no in-depth evaluations of interventions or services, which means that development of models of 'good care' (eg, refs 140 194) still needs to be inferred rather than being based on sound evidence. Further mapping and evaluation of models of care would be beneficial to understand better their differential impact on accessibility, outcomes, cost and patient satisfaction. Finally, there was no research that specifically explored women's/girls' experiences of healthcare professional's involvement in the implementation of legal safeguarding or prevention interventions, which in some countries (such as the UK) are inserting new complexities into the patient–professional relationship and reportedly undermining trust.^{32 33 195–197}

CONCLUSION

This review has identified key challenges but also opportunities for the development of culturally safe and accessible services to improve care for women affected by FGM/C. Future research should involve communities to evaluate existing models of care in order to inform best practice.

Acknowledgements We would like to thank all those who have contributed time and valuable insights to this project. We would particularly like to thank all those who gave up their day to attend the national stakeholder consultation event. Special thanks go to Carol McCormick (FGM/C specialist midwife) for her support and input at the start of this project. We gratefully acknowledge the invaluable contributions of our project advisory group: Kinsi Clarke, Nottingham and Nottinghamshire Refugee Forum, cofounder of Nottingham FGM survivor's group; Professor Jim Thornton, consultant obstetrician, Nottingham University Hospital Trust and University of Nottingham; Amanda Wickham (FGM/C specialist midwife), Nottingham University Hospital Trust; Helen Denness, consultant in public health, Nottingham City Council; and Grace Brough, insight specialist (public health), Nottingham City Council, Chair of Nottingham and Nottinghamshire FGM/C Steering Group.

Contributors CE and VN conceived the study. CE obtained funding, oversaw all aspects of the project and contributed to all stages. CE drafted this paper. RT contributed to all stages of the review, including data extraction, coding, quality appraisal and CERQual components. JE designed and executed all the searches. JM assisted with quality assessment and contributed to the evolving synthesis. GMAH assisted with quality assessment and contributed to the evolving synthesis. JA provided clinical expertise and contributed to the evolving synthesis and formulation of recommendations/conclusions. VN provided a community perspective and contributed to the evolving synthesis and formulation of recommendations/conclusions.

Funding This project was funded by the NIHR HS&DR Programme (no 15/137/04). The views expressed in this report are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given,

and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

REFERENCES

- Zeeb H, Makarova N, Brand T, *et al*. Superdiversity – a new concept for migrant health? *Public Health Forum* 2015;23:124–5.
- Vertovec S. Super-diversity and its implications. *Ethn Racial Stud* 2007;30:1024–54.
- Rechel B, Mladovsky P, Ingleby D, *et al*. Migration and health in an increasingly diverse Europe. *Lancet* 2013;381:1235–45.
- Phillimore J. Approaches to health provision in the age of super-diversity: accessing the NHS in Britain's most diverse city. *Crit Soc Policy* 2011;31:5–29.
- Phillimore J. Migrant maternity in an era of superdiversity: new migrants' access to, and experience of, antenatal care in the West Midlands, UK. *Soc Sci Med* 2016;148:152–9.
- United Nations Children's Fund. *Female Genital Mutilation/Cutting: A Global Concern*. New York: UNICEF, 2016.
- Van Baelen L, Ortensi L, Leye E. Estimates of first-generation women and girls with female genital mutilation in the European Union, Norway and Switzerland. *Eur J Contracept Reprod Health Care* 2016;21:474–82.
- WHO. *Female Genital Mutilation: Fact Sheet*. Geneva: World Health Organisation, 2014.
- WHO. *WHO Guidelines on the Management of Health Complications from Female Genital Mutilation*. Geneva: WHO, 2016.
- Royal College of Obstetricians and Gynaecologists. *Green-Top Guideline No.53: Female Genital Mutilation and its Management*. London: Royal College of Obstetricians and Gynaecologists, 2015.
- Royal College of Midwives, Royal College of Nursing, Royal College of Obstetricians and Gynaecologists, *et al*. *Intercollegiate Guidelines Tackling FGM in the UK: Intercollegiate Recommendations for Identifying, Recording and Reporting*. London 2013.
- Royal College of Nursing, Royal College of Nursing. *Female Genital Mutilation: An RCN Resource for Nursing and Midwifery Practice*. 2nd edn. London, 2015.
- Abdulcadir J, Rodriguez MI, Say L, *et al*. Research gaps in the care of women with female genital mutilation: an analysis. *BJOG* 2015;122:294–303.
- Esu E, Udo A, Okusanya BO, *et al*. Antepartum or intrapartum deinfibulation for childbirth in women with type III female genital mutilation: A systematic review and meta-analysis. *Int J Gynaecol Obstet* 2017;136(Suppl 1):21–9.
- Albert J, Bailey E, Duaso M. Does the timing of deinfibulation for women with type 3 female genital mutilation affect labour outcomes? *Br J Midwifery* 2015;23:430–7.
- Paliwal P, Ali S, Bradshaw S, *et al*. Management of type III female genital mutilation in Birmingham, UK: a retrospective audit. *Midwifery* 2014;30:282–8.
- Okusanya BO, Oduwole O, Nwachuku N, *et al*. Deinfibulation for preventing or treating complications in women living with type III female genital mutilation: A systematic review and meta-analysis. *Int J Gynaecol Obstet* 2017;136(Suppl 1):13–20.
- Gayle C, Rymer J. Female genital mutilation and pregnancy: associated risks. *Br J Nurs* 2016;25:978–83.
- NHS England. *Commissioning Services to Meet the Needs of Women and Girls with FGM*. London 2018.
- WHO. *Care of Girls and Women Living with Female Genital Mutilation: A Clinical Handbook*. Geneva: World Health Organisation, 2018.
- Johansen REB, Ziyada MM, Shell-Duncan B, *et al*. Health sector involvement in the management of female genital mutilation/cutting in 30 countries. *BMC Health Serv Res* 2018;18:240.
- Waigwa S, Doos L, Bradbury-Jones C, *et al*. Effectiveness of health education as an intervention designed to prevent female genital mutilation/cutting (FGM/C): a systematic review. *Reprod Health* 2018;16:62.
- Ahlberg BM, Krantz I, Lindmark G, *et al*. 'It's Only a Tradition': Making Sense of Eradication Interventions and the Persistence of Female 'Circumcision' within a Swedish Context. *Crit Soc Policy* 2004;24:50–78.
- Berg RC, Denison E. A tradition in transition: factors perpetuating and hindering the continuance of female genital mutilation/cutting (FGM/C) summarized in a systematic review. *Health Care Women Int* 2013;34:837–59.
- Recchia N, McGarry J. "Don't judge me": narratives of living with FGM. *Int J Hum Rights Healthc* 2017;10:4–13.
- Baillet H, Murray N, Connelly E, *et al*. Addressing female genital mutilation in Europe: a scoping review of approaches to participation, prevention, protection, and provision of services. *Int J Equity Health* 2018;17:21.
- Dawson A, Homer CS, Turkmani S, *et al*. A systematic review of doctors' experiences and needs to support the care of women with female genital mutilation. *Int J Gynaecol Obstet* 2015;131:35–40.
- Dawson A, Turkmani S, Fray S, *et al*. Evidence to inform education, training and supportive work environments for midwives involved in the care of women with female genital mutilation: a review of global experience. *Midwifery* 2015;31:229–38.
- Reig-Alcaraz M, Siles-Gonzalez J, Solano-Ruiz C. A mixed-method synthesis of knowledge, experiences and attitudes of health professionals to female genital mutilation. *J Adv Nurs* 2015.
- Evans C, Tweheyo R, McGarry J, *et al*. Crossing cultural divides: A qualitative systematic review of factors influencing the provision of healthcare related to female genital mutilation from the perspective of health professionals. *PLoS One* 2019;14:e0211829.
- Connelly E, Murray N, Baillet H, *et al*. Missing from the debate? a qualitative study exploring the role of communities within interventions to address female genital mutilation in Europe. *BMJ Open* 2018;8:e021430.
- Dixon S, Agha K, Ali F, *et al*. Female genital mutilation in the UK—where are we, where do we go next? Involving communities in setting the research agenda. *Res Involv Engagem* 2018;4:29.
- Plugge E, Adam S, El Hindi L, *et al*. The prevention of female genital mutilation in England: what can be done? *J Public Health* 2018;367.
- Evans C, Tweheyo R, McGarry J, *et al*. What are the experiences of seeking, receiving and providing FGM-related healthcare? Perspectives of health professionals and women/girls who have undergone FGM: protocol for a systematic review of qualitative evidence. *BMJ Open* 2017;7:e018170.
- Hamid A, Grace K, Warren N. A meta-synthesis of the birth experiences of African immigrant women affected by female genital cutting. *J Midwifery Womens Health* 2018;00:1–11.
- Turkmani S, Homer CSE, Dawson A. Maternity care experiences and health needs of migrant women from female genital mutilation-practicing countries in high-income contexts: A systematic review and meta-synthesis. *Birth* 2018.
- Smith H, Stein K. Health information interventions for female genital mutilation. *Int J Gynaecol Obstet* 2017;136(Suppl 1):79–82.
- Smith H, Stein K. Surgical or medical interventions for female genital mutilation. *Int J Gynaecol Obstet* 2017;136(Suppl.1):43–6.
- Smith H, Stein K. Psychological and counselling interventions for female genital mutilation. *Int J Gynaecol Obstet* 2017;136(Suppl 1):60–4.
- Jordal M, Wahlberg A. Challenges in providing quality care for women with female genital cutting in Sweden – A literature review. *Sex Reprod Healthc* 2018;17:91–6.
- Scamell M, Ghumman A. The experience of maternity care for migrant women living with female genital mutilation: A qualitative synthesis. *Birth* 2019;46:15–23.
- Suphanchaimat R, Kantamaturapoj K, Putthasri W, *et al*. Challenges in the provision of healthcare services for migrants: a systematic review through providers' lens. *BMC Health Serv Res* 2015;15:390.
- Robertshaw L, Dhesi S, Jones LL. Challenges and facilitators for health professionals providing primary healthcare for refugees and asylum seekers in high-income countries: a systematic review and thematic synthesis of qualitative research. *BMJ Open* 2017;7:e015981.
- O'Donnell CA, Higgins M, Chauhan R, *et al*. Asylum seekers' expectations of and trust in general practice: a qualitative study. *Br J Gen Pract* 2008;58:e1–e11.
- Ochieng BM. Black African migrants: the barriers with accessing and utilizing health promotion services in the UK. *Eur J Public Health* 2013;23:265–9.
- Agudelo-Suárez AA, Gil-González D, Vives-Cases C, *et al*. A metasynthesis of qualitative studies regarding opinions and perceptions about barriers and determinants of health services' accessibility in economic migrants. *BMC Health Serv Res* 2012;12:461.
- Hannes K, Harden A. Multi-context versus context-specific qualitative evidence syntheses: combining the best of both. *Res Synth Methods* 2011;2:271–8.
- Weir MC, Grimshaw JM, Mayhew A, *et al*. Decisions about lumping vs. splitting of the scope of systematic reviews of complex interventions are not well justified: A case study in systematic reviews of health care professional reminders. *J Clin Epidemiol* 2012;65:756–63.

49. Joanna Briggs Institute. JBI Reviewer's Manual. <https://wiki.joannabriggs.org/display/MANUAL/Chapter+2%3A+Systematic+reviews+of+qualitative+evidence> [Accessed 12-07-18].
50. Noyes J, Hannes K, Booth A, et al. Qualitative and Implementation Evidence and Cochrane Reviews, Version 1.02a. In: Higgins JA, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions Version 5/3: The Cochrane Collaboration*, 2015. updated October 2015.
51. Ramsden I. *Cultural Safety and Nursing Education in Aotearoa and Te Waipounamu*. Victoria University 2002.
52. Browne AJ, Varcoe C, Smye V, et al. Cultural safety and the challenges of translating critically oriented knowledge in practice. *Nurs Philos* 2009;10:167–79.
53. Phiri J, Dietsch E, Bonner A. Cultural safety and its importance for Australian midwifery practice. *Collegian* 2010;17:105–11.
54. Duke J, Connor M, McEldowney R. Becoming a culturally competent health practitioner in the delivery of culturally safe care. *J Cult Divers* 2009;16:40–9.
55. Gerlach AJ. A critical reflection on the concept of cultural safety. *Can J Occup Ther* 2012;79:151–8.
56. Cooney C. A comparative analysis of transcultural nursing and cultural safety. *Nurs Pract N Z* 1994;9:6–11.
57. Brascoupé S, Waters C. Cultural safety: exploring the applicability of the concept of cultural safety to aboriginal health and community wellness. *J Aborig Health* 2009;6–41.
58. Wepa D. *Cultural Safety in Aotearoa New Zealand*. North Shore City: Pearson Prentice Hall, 2005.
59. Johnstone MJ, Kanitsaki O. An exploration of the notion and nature of the construct of cultural safety and its applicability to the Australian health care context. *J Transcult Nurs* 2007;18:247–56.
60. Kinn LG, Holgersen H, Ekeland TJ, et al. Metasynthesis and bricolage: an artistic exercise of creating a collage of meaning. *Qual Health Res* 2013;23:1285–92.
61. France EF, Uny I, Ring N, et al. A methodological systematic review of meta-ethnography conduct to articulate the complex analytical phases. *BMC Med Res Methodol* 2019;19:35.
62. Guba E, Lincoln Y. Competing paradigms in qualitative research. In: Denzin N, Lincoln Y, eds. *Handbook of Qualitative Research*. London: Sage, 1994:105–17.
63. Carroll C, Booth A, Leaviss J, et al. “Best fit” framework synthesis: refining the method. *BMC Med Res Methodol* 2013;13.
64. Reeves S, Albert M, Kuper A, et al. Why use theories in qualitative research? *BMJ* 2008;337:a949.
65. Maxwell J. *Qualitative Research Design: An Interactive Approach*. Los Angeles: Sage, 2013.
66. Giacomini M. Theory matters in qualitative health research. In: Bourgeault I, Dingwall R, De Vries R, eds. *The Sage Handbook of Qualitative Methods in Health Research*. London: Sage, 2010:125–55.
67. Oliver S, Dickson K, Bangpan M. *Systematic reviews: making them policy relevant. A briefing for policy makers and systematic reviewers*. London: EPPI-Centre, Social Science Research Unit, UCL Institute of Education, University College London, 2015.
68. Tong A, Flemming K, McInnes E, et al. Enhancing transparency in reporting the synthesis of qualitative research: ENTREQ. *BMC Med Res Methodol* 2012;12:181.
69. Evans C, McGarry J, McCormick C, et al. The experiences of healthcare across the life course for women and girls who have undergone female genital mutilation: a qualitative systematic review Protocol, PROSPERO, CRD42015030001. 2015.
70. National Information Center on Health Services Research and Healthcare Technology (NICHSR). *Health Services Research and Health Policy Grey Literature Project: Summary Report: U. S. National Library of Medicine*, 2006.
71. Adams J, Hillier-Brown FC, Moore HJ, et al. Searching and synthesising ‘grey literature’ and ‘grey information’ in public health: critical reflections on three case studies. *Syst Rev* 2016;5:164.
72. Mahood Q, Van Eerd D, Irvin E. Searching for grey literature for systematic reviews: challenges and benefits. *Res Synth Methods* 2014;5:221–34.
73. Adams RJ, Smart P, Huff AS. Shades of Grey: Guidelines for Working with the Grey Literature in Systematic Reviews for Management and Organizational Studies. *Int J Manag Rev* 2017;19:432–54.
74. Pearson A, Robertson-Mait S, Rittenmeyer L. *Synthesizing Qualitative Evidence*. Philadelphia: Lippincott-Joanna Briggs Institute, 2011.
75. Joanna Briggs Institute. JBI QARI Critical Appraisal Checklist for Interpretive and Critical Research. <https://wiki.joannabriggs.org/display/MANUAL/Joanna+Briggs+Institute+Reviewer%27s+Manual> [Accessed 08-06-18]; cited included reference from Health FGM 170518.enl. Available: http://joannabriggs.org/assets/docs/jbc/operations/criticalAppraisalForms/JBC_Form_CritAp_IntCrit.pdf
76. Munthe-Kaas H, Bohren MA, Glenton C, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 3: how to assess methodological limitations. *Implement Sci* 2018;13:9.
77. Lewin S, Glenton C, Munthe-Kaas H, et al. Using qualitative evidence in decision making for health and social interventions: an approach to assess confidence in findings from qualitative evidence syntheses (GRADE-CERQual). *PLoS Med* 2015;12:e1001895.
78. Dixon-Woods M, Sutton A, Shaw R, et al. Appraising qualitative research for inclusion in systematic reviews: a quantitative and qualitative comparison of three methods. *J Health Serv Res Policy* 2007;12:42–7.
79. Dixon-Woods M, Booth A, Sutton AJ. Synthesizing qualitative research: a review of published reports. *Qual Res* 2007;7:375–422.
80. Dixon-Woods M, et al. The problem of appraising qualitative research. *Qual Saf Health Care* 2004;13:223–5.
81. Sandelowski M, Barroso J. Reading qualitative studies. *Int J Qual Methods* 2002;1:74–108.
82. Noyes J, Booth A, Flemming K, et al. Cochrane Qualitative and Implementation Methods Group guidance paper 3: methods for assessing methodological limitations, data extraction and synthesis, and confidence in synthesized qualitative findings. *J Clin Epidemiol* 2017.
83. Higginbottom GMA, Morgan M, Alexandre M, et al. Immigrant women’s experiences of maternity-care services in Canada: a systematic review using a narrative synthesis. *Syst Rev* 2015;4:1–13.
84. Higginbottom GMA, Hadziabdic E, Yohani S, et al. Immigrant women’s experience of maternity services in Canada: a meta-ethnography. *Midwifery* 2014;30:544–59.
85. Popay J, Rogers A, Williams G. Rationale and standards for the systematic review of qualitative literature in health services research. *Qual Health Res* 1998;8:341–51.
86. Noyes J, Popay J. Directly observed therapy and tuberculosis: how can a systematic review of qualitative research contribute to improving services? A qualitative meta-synthesis. *J Adv Nurs* 2007;57:227–43.
87. Sandelowski M, Barroso J. Finding the findings in qualitative studies. *J Nurs Scholarsh* 2002;34:213–9.
88. Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Med Res Methodol* 2008;8:45.
89. Baldeh F. *Obstetric Care in Scotland: The Experience of Women who have Undergone Female Genital Mutilation (FGM) [MSc Thesis]*. Queen Margaret University, 2013.
90. d’Entremont M, Smythe L, McAra-Couper J. The sounds of silence—a hermeneutic interpretation of childbirth post excision. *Health Care Women Int* 2014;35:300–19.
91. Fawcett L. *Somali Refugee Women and their U.S. Healthcare Providers: Knowledge, Perceptions and Experiences of Childbearing [Doctor of Philosophy]*: Arizona State University, 2014.
92. Glover J, Liebling H, Barrett H, et al. The psychological and social impact of female genital mutilation: a holistic conceptual framework. *J Int Stud* 2017;10:219–38.
93. Jones A. *Working Psychologically with Female Genital Mutilation: An Exploration of the Views and Experiences of Women who have Experienced FGM and of Clinical Psychologists*: University of East London, 2010.
94. Lundberg PC, Gereziher A. Experiences from pregnancy and childbirth related to female genital mutilation among Eritrean immigrant women in Sweden. *Midwifery* 2008;24:214–25.
95. Moxey JM, Jones LL. A qualitative study exploring how Somali women exposed to female genital mutilation experience and perceive antenatal and intrapartum care in England. *BMJ Open* 2016;6:e009846.
96. Palfreyman A, Brown E, Nam S. *Understanding Female Genital Mutilation in Birmingham: Findings from a PEER Study: Options Consultancy Services and Birmingham & Solihull Women’s Aid*, 2011.
97. Straus L, McEwen A, Hussein FM. Somali women’s experience of childbirth in the UK: perspectives from Somali health workers. *Midwifery* 2009;25:181–6.
98. Thierfelder C. *Female Genital Mutilation and the Swiss Health Care System*. University of Basel, 2003.
99. Vangen S, Johansen REB, Sundby J, et al. Qualitative study of perinatal care experiences among Somali women and local health care professionals in Norway. *Eur J Obstet Gynecol Reprod Biol* 2004;112:29–35.

100. Vaughan C, White N, Keogh L, et al. *Listening to North Yarra Communities about Female Genital Cutting*. Melbourne, Australia: The University of Melbourne, 2014.
101. Vaughan C, White N, Keogh L, et al. *Female Genital Mutilation/ Cutting in Regional Victoria. Research to Practice*. Melbourne, Australia: The University of Melbourne, 2014.
102. Boyatzis R. *Transforming Qualitative Data: Thematic Analysis and Code Development*. London: Sage, 1998.
103. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;3:77–101.
104. Booth A, Carroll C, Iltott I, et al. Desperately seeking dissonance: identifying the disconfirming case in qualitative evidence synthesis. *Qual Health Res* 2013;23:126–41.
105. Noyes J, Booth A, Lewin S, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 6: how to assess relevance of the data. *Implement Sci* 2018;13:4.
106. Noyes J, Booth A, Flemming K, et al. Cochrane Qualitative and Implementation Methods Group guidance series—paper 3: methods for assessing methodological limitations, data extraction and synthesis, and confidence in synthesized qualitative findings. *J Clin Epidemiol* 2018;97:49–58.
107. Lewin S, Booth A, Glenton C, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings: introduction to the series. *Implement Sci* 2018;13:2.
108. Lewin S, Bohren M, Rashidian A, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 2: how to make an overall CERQual assessment of confidence and create a Summary of Qualitative Findings table. *Implement Sci* 2018;13:10.
109. Glenton C, Carlsen B, Lewin S, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 5: how to assess adequacy of data. *Implement Sci* 2018;13:14.
110. Colvin CJ, Garside R, Wainwright M, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 4: how to assess coherence. *Implement Sci* 2018;13:13.
111. Booth A, Lewin S, Glenton C, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 7: understanding the potential impacts of dissemination bias. *Implement Sci* 2018;13:12.
112. Asefaw F. *Female Genital Mutilation: A Field Study with Special Consideration of the Background as well as the Health and Psychosexual Consequences for those Affected and their Partners in Eritrea and in Germany* [Weibliche Genitalbeschneidung: Eine Feldstudie Unter besonderer Berücksichtigung der Hintergründe Sowie der Gesundheitlichen und Psychosexuellen Folgen für Betroffene und Partner in Eritrea und Deutschland]. Humboldt University of Berlin, 2007.
113. Maier C. *Echoes of Silence: Voices of Concern for Genital Mutilation in African Immigrant Women in Vienna: An Ethnological Study* [Echo des Schweigens: Stimmen der Betroffenheit zur Genitalverstümmelung bei Afrikanischen Immigrantinnen in Wien: Ethnologische Studie]. University of Vienna, 2003.
114. Ballesteros Meseguer C, Almansa Martinez P, MdM PB, et al. The voice of women subjected to female genital mutilation in the region of Murcia, Spain] La Voz de las mujeres sometidas a mutilación genital femenina en la region de Murcia. *Gac Sanit* 2014;28:287–91.
115. Bravo Pastor del Mar M. *Women Subjected to Female Genital Mutilation: Knowledge for the Nursing Discipline* [La Voz de las Mujeres Sometidas a Mutilación Genital Femenina: Saberes Para la Disciplina Enfermera [Tesis Doctoral]]. University of Murcia, 2014.
116. Guerin PB, Allotey P, Hussein Elmi F, et al. Advocacy as a means to an end: assisting refugee women to take control of their reproductive health needs. *Women Health* 2006;43:7–25.
117. Murray L, Windsor C, Parker E, et al. The experiences of African women giving birth in Brisbane, Australia. *Health Care Women Int* 2010;31:458–72.
118. Betts V. *Transnational Gender, Sexuality and Identity Construction: Ambivalent Subjectivities of Somali Refugee Women in the GTA*. MA Thesis, The University of Guelph, 2011.
119. Chalmers B, Omer-Hashi K. What Somali women say about giving birth in Canada. *J Reprod Infant Psychol* 2002;20:267–82.
120. Glazer E. *Embodiment, Pain and Circumcision in Somali-Canadian Women* [M.Sc.]: University of Toronto, 2012.
121. Khaja K. *Female Circumcision: Life Histories of Somali Women* [Doctor of Philosophy]: The University of Utah, 2004.
122. Khaja K, Lay K, Boys S. Female circumcision: toward an inclusive practice of care. *Health Care Women Int* 2010;31:686–99.
123. Shermarke MAA. *Understanding the Canadian Community Context of Female Circumcision* [M.S.W.]: McGill University, 1996.
124. Degni F, Suominen SB, El Ansari W, et al. Reproductive and maternity health care services in Finland: perceptions and experiences of Somali-born immigrant women. *Ethn Health* 2014;19:348–66.
125. Behrendt A. *Listening to African Voices: Female Genital Mutilation/ Cutting among Immigrants in Hamburg: Knowledge, Attitudes and Practice*. Hamburg, Germany: Plan, 2011.
126. Salad J, Verdonk P, de Boer F, et al. "A Somali girl is Muslim and does not have premarital sex - is vaccination really necessary?" A qualitative study into the perceptions of Somali women in the Netherlands about the prevention of cervical cancer. *Int J Equity Health* 2015;14:1–13.
127. Vloeberghs E, van der Kwaak A, Knipscheer J, et al. Coping and chronic psychosocial consequences of female genital mutilation in The Netherlands. *Ethn Health* 2012;17:677–95.
128. Hussien MA. *Services for Women with Female Genital Mutilation in Christchurch: Perspectives of Women and their Health Providers*: University of Canterbury, 2014.
129. Johansen RE. Undoing female genital cutting: perceptions and experiences of infibulation, defibulation and virginity among Somali and Sudanese migrants in Norway. *Cult Health Sex* 2017;19:528–42.
130. Berggren V, Bergstrom S, Edberg A-K. Being different and vulnerable: experiences of immigrant African women who have been circumcised and sought maternity care in Sweden. *J Transcult Nurs* 2006;17:50–7.
131. Essén B, Johnsdotter S, Hovelius B, et al. Qualitative study of pregnancy and childbirth experiences in Somali women resident in Sweden. *BJOG* 2000;107:1507–12.
132. Wiklund H, Aden AS, Högberg U, et al. Somalis giving birth in Sweden: a challenge to culture and gender specific values and behaviours. *Midwifery* 2000;16:105–15.
133. Thierfelder C, Tanner M, Bodiang CM. Female genital mutilation in the context of migration: experience of African women with the Swiss health care system. *Eur J Public Health* 2005;15:86–90.
134. O'Brien O, Baldeh F, Hassan J, et al. *My Voice: Participatory Action Research Project with Men, Women and Young People on Female Genital Mutilation (FGM) in Scotland: (Phase 2)*. Waverly Care, Edinburgh: Queen Margaret University, 2017.
135. O'Brien O, Baldeh F, Sivapatham S, et al. *Participatory Action Research Project with Men, Women and Young People on Female Genital Mutilation (FGM) in Scotland: (Phase 1)*. Waverly Care, Edinburgh: Queen Margaret University, 2016.
136. Abdi R. Carving culture: Creating identity through female genital cutting. *Durham Anthropology Journal* 2012;18:115–53.
137. Abdullahi A, Copping J, Kessel A, et al. Cervical screening: perceptions and barriers to uptake among Somali women in Camden. *Public Health* 2009;123:680–5.
138. Ahmed M. Attitudes towards FGM among Somali women living in the UK. In: Momoh C, ed. *Female Genital Mutilation*. London: Radcliffe, 2005:95–117.
139. Bulman KH, McCourt C. Somali refugee women's experiences of maternity care in west London: a case study. *Crit Public Health* 2002;12:365–80.
140. Hussein E. *Women's Experiences, Perceptions and Attitudes of Female Genital Mutilation: The Bristol PEER Study*. London: FORWARD, 2010.
141. Maternity Action. *Women's Voices on Health: Addressing Barriers to Accessing Primary Care*. London: Women's Health and Equality Consortium, 2014.
142. Norman K, Gegzabher SB, Otoo-Oyortey N. "Between Two Cultures": A Rapid PEER Study Exploring Migrant Communities' Views on Female Genital Mutilation in Essex and Norfolk, UK. *FORWARD & National FGM Centre Report* 2016.
143. Norman K, Hemmings J, Hussein E, et al. "FGM is Always With Us": Experiences, Perceptions and Beliefs of Women Affected by Female Genital Mutilation in London: Results from a PEER Study. London: Options Consultancy Services and FORWARD, 2009.
144. Safari F. A qualitative study of women's lived experience after defibulation in the UK. *Midwifery* 2013;29:154–8.
145. Ameresekere M, Borg R, Frederick J, et al. Somali immigrant women's perceptions of cesarean delivery and patient-provider communication surrounding female circumcision and childbirth in the USA. *Int J Gynecol Obstet* 2011;115:227–30.
146. Beine K, Fullerton J, Palinkas L, et al. Conceptions of prenatal care among Somali women in San Diego. *J Nurs Midwifery* 1995;40:376–81.
147. Carroll J, Epstein R, Fiscella K, et al. Caring for Somali women: implications for clinician-patient communication. *Patient Educ Couns* 2007;66:337–45.
148. Gali MA. *Female Circumcision: A Transcultural Study of Attitudes, Identity and Reproductive Health of East African Immigrants* [Degree of Doctor of Philosophy in Psychology]: The Wright Institute, University of Berkeley, 1997.

149. Ghebree RG, Sewali B, Osman S, *et al.* Cervical cancer: barriers to screening in the Somali community in Minnesota. *J Immigr Minor Health* 2015;17:722–8.
150. Hill N, Hunt E, Hyrkäs K. Somali immigrant women's health care experiences and beliefs regarding pregnancy and birth in the United States. *J Transcult Nurs* 2012;23:72–81.
151. McNeely S, Christie-de Jong F. Somali refugees' perspectives regarding FGM/C in the US. *Int J Migr Health Soc Care* 2016;12:157–69.
152. Shaw E. Female circumcision: perceptions of clients and caregivers. *J Am Coll Health* 1985;33:193–7.
153. Upvall MJ, Mohammed K, Dodge PD. Perspectives of Somali Bantu refugee women living with circumcision in the United States: a focus group approach. *Int J Nurs Stud* 2009;46:360–8.
154. Lincoln Y, Guba E, *Naturalistic Inquiry*. California: Sage, 1985.
155. Essén B, Binder P, Johnsdotter S. An anthropological analysis of the perspectives of Somali women in the West and their obstetric care providers on caesarean birth. *J Psychosom Obstet Gynaecol* 2011;32:10–18.
156. Green G, Davison C, Bradby H, *et al.* Pathways to care: how superdiversity shapes the need for navigational assistance. *Social Health Illn* 2014;36:1205–19.
157. Phillimore J, Klaas F, Padilla B, *et al.* *Adaptation of Health Services to Diversity: An Overview of Approaches, Working Paper No. 15*: University of Birmingham: Institute for Research into Superdiversity, 2016.
158. Benza S, Liamputtong P. Pregnancy, childbirth and motherhood: a meta-synthesis of the lived experiences of immigrant women. *Midwifery* 2014;30:575–84.
159. Wikberg A, Bondas T. A patient perspective in research on intercultural caring in maternity care: a meta-ethnography. *Int J Qual Stud Health Well-Being* 2010;5:15.
160. Balaam MC, Akerjordet K, Lyberg A, *et al.* A qualitative review of migrant women's perceptions of their needs and experiences related to pregnancy and childbirth. *J Adv Nurs* 2013;69:1919–30.
161. Higginbottom GM, Safipour J, Yohani S, *et al.* An ethnographic study of communication challenges in maternity care for immigrant women in rural Alberta. *Midwifery* 2015;31:297–304.
162. Bohren MA, Vogel JP, Hunter EC, *et al.* The mistreatment of women during childbirth in health facilities globally: a mixed-methods systematic review. *PLoS Med* 2015;12:e1001847.
163. Department of Health. *Midwifery 2020: Delivering Expectations*. London, 2010.
164. NHS England. *National Maternity Review. Better Births: Improving Outcomes of Maternity Services in England*. London, 2016.
165. Mainous AG, Baker R, Love MM, *et al.* Continuity of care and trust in one's physician: evidence from primary care in the United States and the United Kingdom. *Fam Med* 2001;33:22–7.
166. Gilson L. Trust and the development of health care as an institution. *Soc Sci Med* 2003;56:1458–68.
167. Murray B, McCrone S. An integrative review of promoting trust in the patient-primary care provider relationship. *J Adv Nurs* 2015;71:3–23.
168. Denny E. Respectful maternity care needs to be the standard for all women worldwide (mini-commentary). *Br J Obstet Gynaecol* 2017.
169. Shakibazadeh E, Namadian M, Bohren MA, *et al.* Respectful care during childbirth in health facilities globally: a qualitative evidence synthesis. *BJOG* 2018;125.
170. Martin M, Phelan M. Interpreters and cultural mediators: different but complementary roles. *Migr Health* 2010;6:1–8.
171. Akhavan S, Edge D. Foreign-born women's experiences of community based Doulas in Sweden: a qualitative study. *Health Care Women Int* 2012;33:833–48.
172. Akhavan S, Lundgren I. Midwives' experiences of doula support for immigrant women in Sweden: a qualitative study. *Midwifery* 2012;28:80–5.
173. Dundek LH. Establishment of a Somali doula program at a large metropolitan hospital. *J Perinat Neonatal Nurs* 2006;20:128–37.
174. Shelp A. Women helping women: the Somali doula initiative. *Int J Childbirth Educ* 2004;19:4–7.
175. Evans C, Nalubega S, McLuskey J, *et al.* The views and experiences of nurses and midwives in the provision and management of provider-initiated HIV testing and counseling: a systematic review of qualitative evidence. *JBHI Database System Rev Implement Rep* 2016;13:130–286.
176. WHO. *Service Delivery Approaches to HIV Testing and Counselling (HTC): A Strategic Programme Framework*. Geneva: World Health Organisation, 2012.
177. Hamberger LK, Rhodes K, Brown J. Screening and intervention for intimate partner violence in healthcare settings: creating sustainable system-level programs. *J Womens Health* 2015;24:86–91.
178. O'Doherty L, Hegarty K, Ramsay J, *et al.* Screening women for intimate partner violence in healthcare settings. *Cochrane Database Syst Rev* 2015;7(CD007007).
179. Karliner LS, Jacobs EA, Chen AH, *et al.* Do professional interpreters improve clinical care for patients with limited English proficiency? A systematic review of the literature. *Health Serv Res* 2007;42:727–54.
180. Abdulkadir J, Say L, Pallitto C. What do we know about assessing healthcare students and professionals' knowledge, attitude and practice regarding female genital mutilation? A systematic review. *Reprod Health* 2017;14:1–13.
181. Abdulkadir J, Rodriguez MI, Say L. A systematic review of the evidence on clitoral reconstruction after female genital mutilation/cutting. *Int J Gynaecol Obstet* 2015;129:93–7.
182. De Schrijver L, Leye E, Merckx M. A multidisciplinary approach to clitoral reconstruction after female genital mutilation: the crucial role of counselling. *Eur J Contracept Reprod Health Care* 2016;21:269–75.
183. Dominguez Garcia L, Dominguez L, Hopkins K, *et al.* Meeting the sexual health needs of women with female genital mutilation: an integrated model of care. *Int J STD AIDS* 2013;24:2.
184. Low-Beer N, Dominguez-Garcia L, Alcaide D, *et al.* Accessible integrated care to meet the complex needs of women with female genital mutilation (FGM). *Br J Obstet Gynaecol* 2013;120:314.
185. Saha S, Beach MC, Cooper LA. Patient centeredness, cultural competence and healthcare quality. *J Natl Med Assoc* 2008;100:1275–85.
186. Johansen RE. Pain as a counterpoint to culture: toward an analysis of pain associated with infibulation among Somali immigrants in Norway. *Med Anthropol Q* 2002;16:312–40.
187. Johansen RE. Care for infibulated women giving birth in Norway: an anthropological analysis of health workers' management of a medically and culturally unfamiliar issue. *Med Anthropol Q* 2006;20:516–44.
188. James S. Women's experiences of symptoms of posttraumatic stress disorder (PTSD) after traumatic childbirth: a review and critical appraisal. *Arch Womens Ment Health* 2015;18:216–26.
189. Raja S, Hasnain M, Hoersch M, *et al.* Trauma informed care in medicine: current knowledge and future research directions. *Fam Community Health* 2015;38:216–26.
190. Abdulkadir J, Bianchi Demicheli F, Willame A, *et al.* Posttraumatic Stress Disorder Relapse and Clitoral Reconstruction After Female Genital Mutilation. *Obstet Gynecol* 2017;129:371–6.
191. Abayomi O, Chibuzor MT, Okusanya BO, *et al.* Supportive psychotherapy or client education alongside surgical procedures to correct complications of female genital mutilation: A systematic review. *Int J Gynaecol Obstet* 2017;136(Suppl 1):51–5.
192. Adelufosi A, Edet B, Arikpo D, *et al.* Cognitive behavioral therapy for post-traumatic stress disorder, depression, or anxiety disorders in women and girls living with female genital mutilation: A systematic review. *Int J Gynaecol Obstet* 2017;136(Suppl 1):56–9.
193. Jones A, Byrne A. Working psychologically with female genital mutilation: an exploration of the views and experiences of women who have experienced FGM and of clinical psychologists. *Clin Psychol Forum* 2013;246:26–30.
194. Clayton-Hathway K. *A Pilot Evaluation of Health Services for Communities Affected by FGM in Oxfordshire: Final Report*. Oxford, UK: Healthwatch Oxfordshire, 2016.
195. Gangoli G, Gill A, Mulvihill N, *et al.* Perception and barriers: reporting female genital mutilation. *J Aggress Confl Peace Res* 2018.
196. Malik Y, Rowland A, Gerry F, *et al.* Mandatory reporting of female genital mutilation in children in the UK. *Br J Midwifery* 2018;26:377–86.
197. Karlsen S, Carver N, Mogilnicka M, *et al.* *When safeguarding becomes stigmatising: A report on the impact of FGM-safeguarding procedures on people with a Somali heritage living in Bristol, Policy Report*. Bristol: University of Bristol, 2019.